Admissible Diagrams in $U_q^w(\mathfrak{g})$ and Combinatoric Properties of Weyl Groups.

Antoine Mériaux, Gérard Cauchon

Laboratoire d'équations aux dérivées partielles et physique mathématique,
U.F.R. Sciences, B.P. 1039, 51687 Reims Cedex 2, France.

arXiv:0902.0754v1 [math.QA] 4 Feb 2009

Abstract

Consider a complex simple Lie algebra $\mathfrak{g}$ of rank n. Denote by Π a system of simple roots, by W the corresponding Weyl group, consider a reduced expression $w = s_{\alpha_1} \circ \ldots \circ s_{\alpha_t}$ (each $\alpha_i \in \Pi$) of some $w \in W$ and call *diagram* any subset of $[\![1, \ldots, t]\!]$. We denote by $U^+[w]$ (or $U_q^w(\mathfrak{g})$) the "quantum nilpotent" algebra defined as in [11].

We prove (theorem 5.3. 1) that the *positive diagrams* naturally associated with the positive subexpressions (of the reduced expression of w) in the sense of R. Marsh and K. Rietsch [16], coincide with the *admissible diagrams* constructed by G. Cauchon [4] which describe the natural stratification of $Spec(U^+[w])$.

This theorem implies in particular (corollaries 5.3. 1 and 5.3. 2):

1. The map $\zeta : \quad \Delta = \{j_1 < \ldots < j_s\} \mapsto u = s_{\alpha_{j_1}} \circ \ldots \circ s_{\alpha_{j_s}}$ is a bijection from the set of admissible diagrams onto the set $\{u \in W \mid u \leq w\}$.
2. For each admissible diagram $\Delta = \{j_1 < \ldots < j_s\}$, $s_{\alpha_{j_1}} \circ \ldots \circ s_{\alpha_{j_s}}$ is a reduced expression of $u = \zeta(\Delta)$.
3. The map $\zeta' : \quad \Delta = \{j_1 < \ldots < j_s\} \mapsto u' = s_{\alpha_{j_s}} \circ \ldots \circ s_{\alpha_{j_1}}$ is a bijection from the set of admissible diagrams onto the set $\{u \in W \mid u \leq v = w^{-1}\}$.
4. For each admissible diagram $\Delta = \{j_1 < \ldots < j_s\}$, $s_{\alpha_{j_s}} \circ \ldots \circ s_{\alpha_{j_1}}$ is a reduced expression of $u' = \zeta'(\Delta)$.

If the Lie algebra $\mathfrak{g}$ is of type A_n and w is choosen in order that $U^+[w]$ is the quantum matrices algebra $O_q(M_{p,m}(k))$ with $m = n - p + 1$ (see section 2.1), then, by [5], the admissible diagrams are known to be the ⅃ - diagrams in the sense of A. Postnikov [18]. In this particular case, the equality of Le - diagrams and positive subexpressions (of the reduced expression of w) have also been proved (with quite different methods) by A. Postnikov ([18], theorem 19.1.) and by T. Lam and L. Williams ([15], theorem 5.3.).

Contents

1 Introduction. **2**

2 Background on Weyl groups. **4**
2.1 Quantum algebras $U^+[w]$. 4
2.2 Diagrams. 7
2.3 Some properties of positive diagrams. 10

3 Background on the deleting derivations algorithm. **12**
3.1 Conventions. 12
3.2 The algebras $R^{(m)}$. 13
3.3 Prime and H-prime spectrum of $R^{(m)}$, admissible diagrams 15

4 New results on H-invariant prime ideals. **16**
4.1 A necessary and sufficient condition for $\mathcal{P}^{(m)}$ to be in $Im(\phi_m)$. 16
4.2 Some properties of $A^{(m)}$. 18
4.3 Each $x_i^{(m)}$ is a Laurent polynomial in $z_{l_1}, \dots, z_{l_e}$. 21
4.4 A new sufficient condition for $\mathcal{P}^{(m)}$ to be in $Im(\phi_m)$. 26

5 Connections between admissible and positive diagrams. **34**
5.1 A property of admissible diagrams. 34
5.2 The case $w = w_0$. 39
5.3 The general case. 40

1 Introduction.

Given a complex simple Lie algebra $\mathfrak{g}$ of rank n, denote by Π a system of simple roots, by W the corresponding Weyl group, and consider a reduced expression $w = s_{\alpha_1} \circ \dots \circ s_{\alpha_t}$ of some $w \in W$.
Denote by $U^+[w]$ (or $U_q^w(\mathfrak{g})$) the "quantum nilpotent" algebra associated to w as in [11] and by $X_1, \dots, X_t$ the canonical generators of $U^+[w]$ associated with this choosen reduced expression of w (see section 2.1 for more details). The natural action of the torus on the prime spectrum $Spec(U^+[w])$ induces a finite stratification which is completely described by the *admissible diagrams* (also called Cauchon diagrams) which are some particular subsets of $[\![1, \dots, t]\!]$ (see [4]). Those admissible diagrams are very closely connected with the quantum commutation rules satisfied by the canonical generators. For example, in the A_n type, one can choose w such that $U^+[w]$ is a quantum matrices algebra

$$O_q(M_{p,m}(k)) = k < X_{i,j} > \quad (1 \leq i \leq p,\ 1 \leq j \leq m)$$

and we know [5] that the commutation relations

$$X_{u,v}X_{i,j} = X_{i,j}X_{u,v} - (q - q^{-1})X_{i,v}X_{u,j} \quad (i < u,\ j < v) \qquad (*)$$

determine the shape of admissible diagrams by the following observation:
If $\mathcal{P}$ is any completely prime ideal of $O_q(M_{p,m}(k))$, we have immediately

$$X_{u,v} \in \mathcal{P} \Rightarrow (X_{i,v} \in \mathcal{P} \text{ or } X_{u,j} \in \mathcal{P}) \qquad (**)$$

This implies [5] that $\Delta \subset [\![1, \dots, p]\!] \times [\![1, \dots, m]\!]$ (the set of indexes) is an admissible diagram if and only if, for each i, j, u, v as over, we have:

$$(u, v) \in \Delta \Rightarrow ((i, v) \in \mathit{\Delta} \quad \text{or} \quad (u, j) \in \mathit{\Delta})$$

This means that the admissible diagrams are the unions of truncated rows and columns, namely the Ⅎ - diagrams in the sense of A. Postnikov (see [18]).
So, we see on this example that admissible diagrams are quantum objects since, in the non quantum case (when $q = 1$), the formulas $(*)$ become $X_{u,v}X_{i,j} = X_{i,j}X_{u,v}$, so that observation $(**)$ is not valid any longer and the admissible diagrams become invisible.

On the other hand, R. Marsh and K. Rietsch [16] defined the notion of positive subexpression of the reduced w's expression considered over. These positive subexpressions are defined by particular subsets of $[\![1, \dots, t]\!]$ that we call the *positive diagrams.* R. Marsh and K. Rietsch proved in [16] that they are in one to one natural correspondence with the elements of the Weyl group which are smaller or equal to w (for the Bruhat order). In this paper, we prove (theorem 5.3. 1) that *the positive diagrams coincide with the admissible diagrams,* which can be interpreted saying that *R. Marsh and K. Rietsch positive subexpressions are quantum objects.*

In particular, this implies (corollary 5.3. 1):

1. The map $\zeta : \quad \Delta = \{j_1 < \dots < j_s\} \mapsto u = s_{\alpha_{j_1}} \circ \dots \circ s_{\alpha_{j_s}}$ is a bijection from the set of admissible diagrams onto the set $\{u \in W \mid u \leq w\}$,
2. Consider an admissible diagram $\Delta = \{j_1 < \dots < j_s\}$ and some integer $i \in [\![1, \dots, t]\!]$. Set $\Delta \cap [\![i+1, \dots, t]\!] = \{j_c, \dots, j_s\}$ $(1 \leq c \leq s)$. Then the expression $s_{\alpha_i} \circ s_{\alpha_{j_c}} \circ \dots \circ s_{\alpha_{j_s}}$ is reduced. In particular, $s_{\alpha_{j_1}} \circ \dots \circ s_{\alpha_{j_s}}$ is a reduced expression of $\zeta(\Delta)$,
3. Consider some $u \in W$ with $u \leq w$. Then, the only admissible diagram Δ such that $\zeta(\Delta) = u$ is recursively defined as follows:
   - $1 \in \Delta \Leftrightarrow l(s_{\alpha_1} \circ u) = l(u) - 1 \Leftrightarrow u^{-1}(\alpha_1)$ is a negative root,
   - Consider some integer $i \in [\![1, \dots, t-1]\!]$, assume that $\Delta \cap [\![1, \dots, i]\!] = \{j_1, \dots, j_d\}$ and set $u_i = s_{\alpha_{j_d}} \circ \dots \circ s_{\alpha_{j_2}} \circ s_{\alpha_{j_1}} \circ u$ $(u_i = u$ if $\Delta \cap [\![1, \dots, i]\!] = \emptyset)$.

   Then $(i+1 \in \Delta) \Leftrightarrow (l(s_{\alpha_{i+1}} \circ u_i) = l(u_i) - 1) \Leftrightarrow (u_i^{-1}(\alpha_{i+1})$ is a negative root).

It also implies (corollary 5.3. 2):

1. The map $\zeta' : \quad \Delta = \{j_1 < \dots < j_s\} \mapsto u' = s_{\alpha_{j_s}} \circ \dots \circ s_{\alpha_{j_1}}$ is a bijection from the set of admissible diagrams onto the set $\{u \in W \mid u \leq v = w^{-1}\}$.
2. Consider an admissible diagram $\Delta = \{j_1 < \dots < j_s\}$ and some integer $i \in [\![1, \dots, t]\!]$. Set $\Delta \cap [\![i+1, \dots, t]\!] = \{j_c, \dots, j_s\}$ $(1 \leq c \leq s)$. Then the expression $s_{\alpha_{j_s}} \circ \dots \circ s_{\alpha_{j_c}} \circ s_{\alpha_i}$ is reduced. In particular, $s_{\alpha_{j_s}} \circ \dots \circ s_{\alpha_{j_1}}$ is a reduced expression of $\zeta'(\Delta)$.

In the particular case $U^+[w] = O_q(M_{p,m}(k))$ mentioned over, this last result gives back a theorem of A. Postnikov [18] and a theorem of T. Lam and L. Williams [15].

In the general case, let us denote by $\Delta \leftrightarrow w^\Delta$ the one to one correspondence constructed by R. Marsh and K. Rietsch [16] between the positive diagrams and the elements of the Weyl group which are smaller or equal to w.
Let us also denote by $\Delta \leftrightarrow \mathcal{P}_\Delta$ the one to one correspondence constructed by G. Cauchon [4] between the admissible diagrams and the prime ideals of $U^+[w]$ which are invariant under the torus action.
By the theorem 5.3. 1 of this paper, it turns out that there exists a natural one to one correspondence

$$w^\Delta \leftrightarrow \mathcal{P}_\Delta$$

between the elements of the Weyl group which are smaller or equal to w and the prime ideals of $U^+[w]$ which are invariant under the torus action.
Let us recall that, in the particular case $U^+[w] = O_q(M_{p,m}(k))$ mentioned over, S. Launois [13] has constructed (with quite different methods) such a one to one correspondence which, moreover, preserves the ordering (where the Weyl group is provided with the Bruhat order and the spectrum $Spec(O_q(M_{p,m}(k)))$ is provided with the inclusion order).
This leads us to ask the following questions (unsolved at the moment):

- *Question 1.* Does this Launois correspondence coincide with ours in the particular case $U^+[w] = O_q(M_{p,m}(k))$?

- *Question 2.* Does our correspondence $w^\Delta \leftrightarrow \mathcal{P}_\Delta$ preserve the ordering in the general case (where the Weyl group is provided with the Bruhat order and the spectrum $Spec(U^+[w])$ is provided with the inclusion order)?

(A positive answer would supply the Bruhat order with a nice quantum interpretation.)

2 Background on Weyl groups.

Following J. C. Jantzen ([11]), we use the following conventions all along this paper.

- $\mathfrak{g}$ is a complex simple Lie algebra.
- Φ is the (irreducible) root system of $\mathfrak{g}$ with respect to a fixed Cartan subalgebra.
- Π is a fixed basis of Φ, Φ^+ denotes the subset of positive roots, and we set $n = |\Pi|$, $\mathrm{N} = |\Phi^+|$ $(1 \leq n \leq N)$.
- W is the Weyl group of Φ and $(\ ,\ )$ is the unique scalar product on the real vector space V generated by Φ, such that $\|\beta\|^2 = 2$ $(\|\beta\| := \sqrt{(\beta, \beta)})$ for all short roots β in Φ.
- For any β in Φ, we set $d_\beta = \dfrac{\|\beta\|^2}{2}$, $\beta^\vee = \dfrac{\beta}{d_\beta}$, and s_β denotes the reflection with respect to β $(s_\beta(x) = x - (\beta^\vee, x)\beta$ for any x in $V)$.
- k is a field with $\mathrm{char}(k) \neq 2$ and, in addition, $\mathrm{char}(k) \neq 3$ if Φ has type G_2.
- $q \in k^* := k \setminus \{0\}$, q is not a root of unity.
- The k-algebra $U_q(\mathfrak{g})$ and it's canonical generators E_α, F_α, $K_\alpha^{\pm 1}$ $(\alpha \in \Pi)$ are defined as in [11]. We denote by U^+ (or $U_q^+(\mathfrak{g})$) the subalgebra generated by the E_α $(\alpha \in \Pi)$.
- Denote by $\mathbb{Z}\Pi$ the root lattice. The algebra $U_q(\mathfrak{g})$ is $\mathbb{Z}\Pi$-graded and, if α in Π, $\deg(E_\alpha) = \alpha$, $\deg(F_\alpha) = -\alpha$, $\deg(K_\alpha^{\pm 1}) = 0$.
- For any $\rho = \sum_{\alpha \in \Pi} m_\alpha \alpha$ $(m_\alpha \in \mathbb{Z})$ in the root lattice $\mathbb{Z}\Pi$, we set $K_\rho = \prod_{\alpha \in \Pi} K_\alpha^{m_\alpha}$. The multiplicative group $T = \{K_\rho \mid \rho \in \mathbb{Z}\Pi\}$ is called the Torus (of $U_q(\mathfrak{g})$). This group acts on the algebra $U_q(\mathfrak{g})$ by
$$K_\rho.u = K_\rho^{-1} u K_\rho \quad (\forall u \in U_q(\mathfrak{g}))$$
In particular, for any homogeneous element u of $U_q(\mathfrak{g})$ with degree $\deg(u) = \gamma \in \mathbb{Z}\Pi$, we have $K_\rho.u = q^{-(\rho, \gamma)} u$.

2.1 Quantum algebras $U^+[w]$.

Consider any $w \in W$, set $t = l(w)$ and consider a reduced expression
$$w = s_{\alpha_1} \circ \ \dots \ \circ s_{\alpha_t} \qquad (\alpha_i \in \Pi \ \ for \ \ 1 \leq i \leq t) \tag{1}$$
It is well known that
$$\beta_1 = \alpha_1, \ \beta_2 = s_{\alpha_1}(\alpha_2), \ \dots, \ \beta_t = s_{\alpha_1} \dots s_{\alpha_{t-1}}(\alpha_t)$$
are distinct positive roots and that the set $\{\beta_1, \dots, \beta_t\}$ does not depend on the reduced expression (1) of w. For any $\alpha \in \Pi$, define the braid automorphism T_α of the algebra $U_q(\mathfrak{g})$ as in ([11], p. 153), and set
$$X_{\beta_1} = E_{\alpha_1}, \ X_{\beta_2} = T_{\alpha_1}(E_{\alpha_2}), \ \dots, \ X_{\beta_t} = T_{\alpha_1} \dots T_{\alpha_{t-1}}(E_{\alpha_t}).$$
The following results are classical ([11], chapter 8):

- $X_{\beta_1}, \dots, X_{\beta_t}$ are all in U^+. Each X_{β_i} $(1 \leq i \leq t)$ is homogeneous of degree $\deg(X_{\beta_i}) = \beta_i$.
- We denote by $U^+[w]$ (or $U_q^w(\mathfrak{g})$) the subalgebra of U^+ generated by $X_{\beta_1}, \dots, X_{\beta_t}$. This algebra does not depend on the above reduced expression (1) of w (although the variables $X_{\beta_1}, \dots, X_{\beta_t}$ depend of (1)).
- The ordered monomials $X^{\underline{a}} := X_{\beta_1}^{a_1} \dots X_{\beta_t}^{a_t}$, $\underline{a} = (a_1, \dots, a_1) \in \mathbb{N}^t$, are a basis of $U^+[w]$ (as a k-vector space).
- Since the above generators X_{β_i} are homogeneous, the $\mathbb{Z}\Pi$-graduation of $U_q(\mathfrak{g})$ induces a $\mathbb{Z}\Pi$-graduation of $U^+[w]$ and the action of the Torus T on $U_q(\mathfrak{g})$ induces, by restriction, an action of T on $U^+[w]$.

- If $1 \le i < j \le t$, we have the following straightening formula due to Levendorskii and Soibelman:

$$X_{\beta_j} X_{\beta_i} - q^{-(\beta_i, \beta_j)} X_{\beta_i} X_{\beta_j} \ = \ P_{j,i}, \tag{2}$$

$$P_{j,i} \ = \sum_{\underline{a} \ = \ (a_{i+1}, \ldots, a_{j-1})} c_{\underline{a}} X_{\beta_{i+1}}^{a_{i+1}} \ldots X_{\beta_{j-1}}^{a_{j-1}}. \tag{3}$$

  with $\underline{a} \in \mathbb{N}^{j-i-1}$, $c_{\underline{a}} \in k$, and $c_{\underline{a}} \neq 0$ for only finitely many $\underline{a}$. $P_{j,i}$ is homogeneous with degree $\beta_i + \beta_j$ so that, if $j = i+1$, we have $P_{j,i} = 0$.

  The reader will observe a little difference between those formulas and the original Levendorskii - Soibeman's one ([14], prop. 5.5.2.), in which the left member of (2) is $X_{\beta_j} X_{\beta_i} - q^{(\beta_i, \beta_j)} X_{\beta_i} X_{\beta_j}$. The reason is that Levendorskii and Soibelman use a version of the quantum group $U_q(\mathfrak{g})$ which slightly differs from our's. Under our conventions, a direct proof of formulas (2) and (3) is given in [17].

- When $w = w_0$, we have $t = N$, $\{\beta_1, \ldots, \beta_t\} = \Phi^+$ and $U^+[w] = U^+$.

An example

Assume, for sake of simplicity, that $k = \mathbb{C}$ is the complex numbers field, that $\mathfrak{g}$ has type A_n with $n \ge 3$ and that the simple roots $\epsilon_1, \ldots, \epsilon_n$ are ordered such as the Dynkin diagram is

$$\epsilon_1 \ \text{——} \ \epsilon_2 \ \text{——} \ \epsilon_3 \ \text{——} \quad \cdots \quad \text{——} \ \epsilon_{n-1} \ \text{——} \ \epsilon_n$$

Consider the following particular reduced expression of the longest element in W:

$$w_0 \ = s_{\epsilon_1} \circ (s_{\epsilon_2} \circ s_{\epsilon_1}) \circ \ \ldots \ \circ (s_{\epsilon_j} \circ s_{\epsilon_{j-1}} \circ \ \ldots \ \circ s_{\epsilon_1}) \circ \ \ldots \ \circ (s_{\epsilon_n} \circ s_{\epsilon_{n-1}} \circ \ \ldots \ \circ s_{\epsilon_1}) \tag{4}$$

Denote by $Y_{1,2}, Y_{1,3}, Y_{2,3}, \ldots, Y_{1,j+1}, Y_{2,j+1}, \ldots, Y_{j,j+1}, \ldots, Y_{1,n+1}, Y_{2,n+1}, \ldots, Y_{n,n+1}$ the canonical generators of U^+ with respect to this reduced decomposition and observe that:

Lemma 2.1. 1

1. *$Y_{j,j+1} = E_{\alpha_j}$ for $1 \le j \le n$.*
2. *$Y_{i,j+1} = Y_{i,j} Y_{j,j+1} - q^{-1} Y_{j,j+1} Y_{i,j}$ for $1 \le i < j \le n$.*

Proof

1. This results from the equality

$$s_{\epsilon_1} \circ (s_{\epsilon_2} \circ s_{\epsilon_1}) \circ \ \ldots \ \circ (s_{\epsilon_j} \circ s_{\epsilon_{j-1}} \circ \ \ldots \ \circ s_{\epsilon_2})(\epsilon_1) \ = \ \epsilon_j.$$

2. Recall (see section 2.1) that

$$Y_{i,j+1} \ = \ T_{\epsilon_1} \circ (T_{\epsilon_2} \circ T_{\epsilon_1}) \circ \ \ldots \ \circ (T_{\epsilon_{j-1}} \circ T_{\epsilon_{j-2}} \circ \ \ldots \ \circ T_{\epsilon_1}) \circ (T_{\epsilon_j} \circ T_{\epsilon_{j-1}} \circ \ \ldots \ \circ T_{\epsilon_{j+2-i}})(E_{\epsilon_{j+1-i}}).$$

   Since $\{\epsilon_{j-i}, \ldots, \epsilon_1\} \perp \{\epsilon_j, \ldots, \epsilon_{j+2-i}\}$ and $\{\epsilon_{j-i-1}, \ldots, \epsilon_1\} \perp \epsilon_{j+1-i}$, $Y_{i,j+1}$ is equal to

$$T_{\epsilon_1} \circ (T_{\epsilon_2} \circ T_{\epsilon_1}) \circ \ \ldots \ \circ (T_{\epsilon_{j-1}} \circ T_{\epsilon_{j-2}} \circ \ \ldots \ \circ T_{\epsilon_{j+1-i}}) \circ (T_{\epsilon_j} \circ T_{\epsilon_{j-1}} \circ \ \ldots \ \circ T_{\epsilon_{j+2-i}}) \circ T_{\epsilon_{j-i}}(E_{\epsilon_{j+1-i}})$$

   with, by [11],

$$T_{\epsilon_{j-i}}(E_{\epsilon_{j+1-i}}) \ = \ E_{\epsilon_{j-i}} E_{\epsilon_{j+1-i}} \ - \ q^{-1} E_{\epsilon_{j+1-i}} E_{\epsilon_{j-i}}.$$

   Moreover, we have

$$\begin{aligned} & s_{\epsilon_1} \circ (s_{\epsilon_2} \circ s_{\epsilon_1}) \circ \ \ldots \ \circ (s_{\epsilon_{j-1}} \circ s_{\epsilon_{j-2}} \circ \ \ldots \ \circ s_{\epsilon_{j+1-i}}) \circ (s_{\epsilon_j} \circ s_{\epsilon_{j-1}} \circ \ \ldots \ \circ s_{\epsilon_{j+2-i}})(\epsilon_{j+1-i}) \\ & = \ s_{\epsilon_1} \circ (s_{\epsilon_2} \circ s_{\epsilon_1}) \circ \ \ldots \ \circ (s_{\epsilon_{j-1}} \circ s_{\epsilon_{j-2}} \circ \ \ldots \ \circ s_{\epsilon_{j+1-i}})(\epsilon_{j+1-i} + \epsilon_{j+2-i} + \ \ldots \ + \epsilon_j) \\ & = \ s_{\epsilon_1} \circ (s_{\epsilon_2} \circ s_{\epsilon_1}) \circ \ \ldots \ \circ (s_{\epsilon_{j-2}} \circ s_{\epsilon_{j-3}} \circ \ \ldots \ \circ s_{\epsilon_1})(\epsilon_j) \ = \ \epsilon_j. \end{aligned}$$

This implies that

$$T_{\epsilon_1} \circ (T_{\epsilon_2} \circ T_{\epsilon_1}) \circ \ ... \ \circ (T_{\epsilon_{j-1}} \circ T_{\epsilon_{j-2}} \circ \ ... \ \circ T_{\epsilon_{j+1-i}}) \circ (T_{\epsilon_j} \circ T_{\epsilon_{j-1}} \circ \ ... \ \circ T_{\epsilon_{j+2-i}})(E_{\epsilon_{j+1-i}})$$

$$= \ E_{\epsilon_j} \ = \ Y_{j,j+1}.$$

Since $\{\epsilon_j, \ \epsilon_{j-1} \ ... \ , \epsilon_{j+2-i}\} \perp \epsilon_{j-i},$ we have

$$T_{\epsilon_1} \circ (T_{\epsilon_2} \circ T_{\epsilon_1}) \circ \ ... \ \circ (T_{\epsilon_{j-1}} \circ T_{\epsilon_{j-2}} \circ \ ... \ \circ T_{\epsilon_{j+1-i}}) \circ (T_{\epsilon_j} \circ T_{\epsilon_{j-1}} \circ \ ... \ \circ T_{\epsilon_{j+2-i}})(E_{\epsilon_{j-i}})$$
$$= \ T_{\epsilon_1} \circ (T_{\epsilon_2} \circ T_{\epsilon_1}) \circ \ ... \ \circ (T_{\epsilon_{j-1}} \circ T_{\epsilon_{j-2}} \circ \ ... \ \circ T_{\epsilon_{j+1-i}})(E_{\epsilon_{j-i}}) \ = \ Y_{i,j}.$$

□

Denote by v a square root of q and, as in [1], denote by $e_{i,j}$ the generators of U^+ constructed by H. Yamane [19]. We know that:

- $e_{j,j+1} = E_{\epsilon_j}$ for $1 \leq j \leq n.$
- $e_{i,j+1} = ve_{i,j}e_{j,j+1} - v^{-1}e_{j,j+1}e_{i,j}$ for $1 \leq i < j \leq n.$

This implies:

- $v^{i-j}e_{i,j+1} = (v^{i-j+1}e_{i,j})e_{j,j+1} - e_{j,j+1}q^{-1}(v^{i-j+1}e_{i,j})$ for $1 \leq i < j \leq n$ and, from lemma 2.1. 1, we deduce (by induction on $j-i$):

Lemma 2.1. 2
$Y_{i,j+1} = v^{i-j}e_{i,j+1}$ *for* $1 \leq i \leq j \leq n.$

Now, consider an integer p with $1 < p < n$ and set

$$w \ = \ (s_{\epsilon_p} \circ s_{\epsilon_{p-1}} \circ \ ... \ \circ s_{\epsilon_1}) \circ (s_{\epsilon_{p+1}} \circ s_{\epsilon_p} \circ \ ... \ \circ s_{\epsilon_2}) \circ \ ... \ \circ (s_{\epsilon_n} \circ s_{\epsilon_{n-1}} \circ \ ... \ \circ s_{\epsilon_{n-p+1}}) \quad (5)$$

If $p < j < n,$ we observe that
$\{\epsilon_{j-p}, \ ... \ , \ \epsilon_1\} \perp \{\epsilon_{j+1}, \ ... \ , \ \epsilon_{j+1-p+1}, \ ... \ , \ \epsilon_n, \ ... \ , \ \epsilon_{n-p+1}\}.$
This implies that

$$w_0 \ = \ w_1 \circ w \circ w_2$$

with

$$w_1 \ = s_{\epsilon_1} \circ (s_{\epsilon_2} \circ s_{\epsilon_1}) \circ \ ... \ \circ (s_{\epsilon_{p-1}} \circ s_{\epsilon_{p-2}} \circ \ ... \ \circ s_{\epsilon_1}),$$

$$w_2 \ = s_{\epsilon_1} \circ (s_{\epsilon_2} \circ s_{\epsilon_1}) \circ \ ... \ \circ (s_{\epsilon_{n-p}} \circ s_{\epsilon_{n-p-1}} \circ \ ... \ \circ s_{\epsilon_1}).$$

If $d, \ d_1, \ d_2$ denote the number of simple reflections which appear respectively in the expressions of $w, \ w_1, \ w_2$ given above, we observe that $d + d_1 + d_2$ is equal to the length of $w_0.$ This implies:

Lemma 2.1. 3

1. (5) *is a reduced expression of* $w.$
2. *If* $p \leq j \leq n,$ *then*

$$T_{w_1} \circ (T_{\epsilon_p} \circ T_{\epsilon_{p-1}} \circ \ ... \ \circ T_{\epsilon_1}) \circ \ ... \ \circ (T_{\epsilon_{j-1}} \circ T_{\epsilon_{j-2}} \circ \ ... \ \circ T_{\epsilon_{j-p+1}})(E_{\epsilon_j}) \ = \ Y_{j,j+1}.$$

3. *If* $p \leq j \leq n$ *and* $1 \leq i \leq p,$ *then*

$$T_{w_1} \circ (T_{\epsilon_p} \circ T_{\epsilon_{p-1}} \circ \ ... \ \circ T_{\epsilon_1}) \circ \ ... \ \circ (T_{\epsilon_j} \circ T_{\epsilon_{j-1}} \circ \ ... \ \circ T_{\epsilon_{j-i+2}})(E_{\epsilon_{j-i+1}}) \ = \ Y_{i,j+1}.$$

This implies that the canonical generators $X_{i,j+1}$ of $U^+[w]$ verify

$$X_{i,j+1} \ = \ T_{w_1}^{-1}(Y_{i,j+1})$$

for $p \leq j \leq n$ and $1 \leq i \leq p$.

Since the Yamane generators $e_{i,j+1}$ $(p \leq j \leq n \ \ , \ 1 \leq i \leq p)$ verify the commutation relations of quantum matrices, it results from lemma 2.1. 2 that the generators $Y_{i,j+1}$ verify the same property. Now, as $T_{w_1}^{-1}$ is an automorphism of $U_q(\mathfrak{g})$ and as the ordered monomials in the variables $X_{i,j+1}$ (where the order on these variables is defined by the inverse lexicographic order on the indexes $(i, j+1)$) are a basis of $U^+[w]$, we conclude that

Proposition 2.1. 1
$U^+[w]$ is the algebra $O_q(M_{p,m}(k))$ with $m = n-p+1$ and with canonical generators the variables $X_{i,j+1}$ $(p \leq j \leq n \ \ , \ 1 \leq i \leq p)$.

2.2 Diagrams.

w, t, the simple roots α_i, the positive roots β_i $(1 \leq i \leq t)$, are defined as in section **2.1.,** and we set

$$v \ = \ w^{-1} \ = \ s_{\alpha_t} \circ \ ... \ \circ s_{\alpha_1} \tag{6}$$

Definition 2.2. 1 *A diagram with respect to (1) is any subset Δ of $[\![1, \ ... \ ,t]\!]$. (If there is no possible ambiguity, we omit to precise "with respect to (1)".)*

In the following, we sometimes also omit the symbol $\circ$ in the composition of maps.

Consider a diagram Δ. For any $i \in [\![1, \ ... \ , t]\!]$, we set

$$s_{\alpha_i}^{\Delta} = \left\{ \begin{array}{cc} s_{\alpha_i} & if \ i \ \in \ \Delta \\ Id & if \ i \ \notin \ \Delta \end{array} \right\}$$

and we denote

- $w^{\Delta} \ = \ s_{\alpha_1}^{\Delta} \ ... \ s_{\alpha_t}^{\Delta}$
- $v^{\Delta} \ = \ (w^{\Delta})^{-1} \ = \ s_{\alpha_t}^{\Delta} \ ... \ s_{\alpha_1}^{\Delta}$
- for any $i \in [\![1, \ ... \ , t, t+1]\!]$, $w_i^{\Delta} \ = \ s_{\alpha_i}^{\Delta} \ ... \ s_{\alpha_t}^{\Delta}$ $\quad (w_1^{\Delta} \ = w^{\Delta} \ , w_{t+1}^{\Delta} = Id)$
- for any $i \in [\![0, \ 1, \ ... \ , t]\!]$, $v_i^{\Delta} \ = \ (w_{t-i+1}^{\Delta})^{-1} \ = \ s_{\alpha_t}^{\Delta} \ ... \ s_{\alpha_{t-i+2}}^{\Delta} s_{\alpha_{t-i+1}}^{\Delta}$ $\quad (v_0^{\Delta} \ = \ (w_{t+1}^{\Delta})^{-1} = Id, \ v_t^{\Delta} \ = \ v^{\Delta})$

Remark (The case of example 2.1)

Assume we are in the situation of example 2.1, so that

$$\begin{aligned} w \ &= s_{\alpha_1} \circ s_{\alpha_2} \circ ... \circ s_{\alpha_t} \\ &= (s_{\epsilon_p} \circ s_{\epsilon_{p-1}} \circ \ ... \ \circ s_{\epsilon_1}) \circ (s_{\epsilon_{p+1}} \circ s_{\epsilon_p} \circ \ ... \ \circ s_{\epsilon_2}) \circ \ ... \ \circ (s_{\epsilon_n} \circ s_{\epsilon_{n-1}} \circ \ ... \ \circ s_{\epsilon_{n-p+1}}) \end{aligned}$$

Set $m = n-p+1$, so that $t = mp$, and consider a rectangular tableau consisting in $p \times m$ boxes labeled from 1 to mp as mentioned in the following figure.



Following A. Postnikov [18], *let us draw this tableau with* p *wires going along the rows and* m *wires going along the columns. Label ends of the wires from* 1 *to* $n+1$ *as in the following figure.*



We observe that, if W *is identified as usual with the symmetric group* $\mathcal{S}_{n+1}$ *so that each* s_{ϵ_i} *is the transposition* $(i, i+1)$, *the wiring diagram defined over corresponds to the permutation* $v = w^{-1}$.
Now, consider any diagram Δ (*i.e. any subset of* $[\![1, mp]\!]$), *color in dark the boxes which label is in* Δ, *and replace the corresponding crossing in non-colored boxes as a non crossing :*



white box



colored box

Then, if $\Delta = \{i_1 < ... < i_l\}$, *we observe that this new wiring diagram corresponds to the permutation* $s_{\epsilon_{i_l}} \circ ... \circ s_{\epsilon_{i_1}} = v^{\Delta}$.

So, using the *A. Postnikov's conventions* ([18], p. 71-72) we can say:

1. Δ is a pipe dream or a wiring diagram.
2. $u = v^{\Delta}$ is the element of W corresponding to the pipe dream Δ and Δ is a pipe dream (or a wiring diagram) of u.

   In particular, we have:
3. v is the element of W corresponding to the full pipe dream $\Delta = [\![1, \dots, t]\!]$.
4. Id is the element of W corresponding to the empty pipe dream $\Delta = \emptyset$.

Now, in the general case, let us denote by $\leq$ the Bruhat order on the Weyl group W. Since (1) is a reduced expression of w, we have that (6) is a reduced expression of v. Let us recall ([12], corollary A.1.8) that an element u of W satisfies $u \leq w$ (resp. $u \leq v$) if and only if it can we written as a product of simple refections obtained by omitting some of the s_{α_i} in the reduced expression (1) (resp. (6)). So, we have immediately:

Lemma 2.2. 1

1. *The map:* $\Delta \mapsto u = w^{\Delta}$ *(resp.* $u = v^{\Delta}$*) is surjective from the set of diagrams onto the set* $\{u \in W \mid u \leq w\}$ *(resp.* $\{u \in W \mid u \leq v\}$*).*
2. *The map* $u \mapsto u^{-1}$ *is a bijection from* $\{u \in W \mid u \leq w\}$ *onto* $\{u \in W \mid u \leq v\}$*.*

Lemma 2.2. 2
$\mathbf{v}^{\Delta} := (v_0^{\Delta}, v_1^{\Delta}, \dots, v_t^{\Delta})$ is a subexpression of (6) in the sense of Marsh and Rietsch [16].

Proof

This is because $v_{i-1}^{\Delta} = s_{\alpha_t}^{\Delta} \dots s_{\alpha_{t-i+2}}^{\Delta} \implies v_i^{\Delta} = v_{i-1}^{\Delta} s_{\alpha_{t-i+1}}^{\Delta}$ with

$$s_{\alpha_{t-i+1}}^{\Delta} = \left\{ \begin{array}{cc} s_{\alpha_{t-i+1}} & if \ \ t-i+1 \in \Delta \\ Id & if \ \ t-i+1 \notin \Delta \end{array} \right\}.$$

$\square$

Lemma 2.2. 3
If $\mathbf{u} = (u_0, u_1, \dots, u_t)$ *is a subexpression of* v*, there exists a unique diagram* Δ *such that* $\mathbf{v}^{\Delta} = \mathbf{u}$ *. So, the map* $\Delta \mapsto \mathbf{v}^{\Delta}$ *is a bijection from the set of diagrams onto the set of subexpressions of* v*.*

Proof

Since $\mathbf{u}$ is a subexpression of v, we have $u_0 = Id$ and

$$(\forall i \in [\![1, \dots, t]\!]) \quad (u_{i-1})^{-1} u_i \in \{s_{t-i+1}, \ Id\}.$$

Given any diagram Δ, we have $\mathbf{v}^{\Delta} = (v_0^{\Delta}, v_1^{\Delta}, \dots, v_t^{\Delta})$. Since $v_0^{\Delta} = u_0 = Id$, we have $\mathbf{v}^{\Delta} = \mathbf{u}$ if and only if

$$(\forall i \in [\![1, \dots, t]\!]) \quad (u_{i-1})^{-1} u_i = (v_{i-1}^{\Delta})^{-1} v_i^{\Delta} = s_{\alpha_{t-i+1}}^{\Delta}.$$

Now, there exists a unique diagram Δ which satisfies those conditions. It is defined by

$$(\forall i \in [\![1, \dots, t]\!]) \quad (t-i+1 \in \Delta) \Leftrightarrow (u_{i-1})^{-1} u_i = s_{t-i+1}.$$

$\square$

Definition 2.2. 2 *We say that a diagram* Δ *is positive with respect to the reduced decomposition (1) (or that* Δ *is positive if there is no ambiguity) if* $\mathbf{v}^{\Delta} = (v_0^{\Delta}, v_1^{\Delta}, \dots, v_t^{\Delta})$ *is a positive subexpression of (6) in the sense of Marsh and Rietsch* [16]*, namely:*

$$v_{i-1}^{\Delta} < v_{i-1}^{\Delta} s_{\alpha_{t-i+1}} \quad (\forall \ i \in [\![1, \dots, t]\!])$$

Observation 2.2. 1 *The full diagram and the empty diagram are both positive.*

Proof

- If $\Delta = [\![1, \dots ,t]\!]$ is the full diagram, we have $v^{\Delta}_{i-1} = s_{\alpha_t} \dots s_{\alpha_{t-i+2}}$ for any $i \in [\![1, \dots ,t]\!]$ $(v^{\Delta}_0 = Id)$. Since (6) is a reduced expression of v, $s_{\alpha_t} \dots s_{\alpha_{t-i+2}} s_{\alpha_{t-i+1}}$ is a reduced expression of $v^{\Delta}_{i-1} s_{\alpha_{t-i+1}}$. So, clearly, we have $v^{\Delta}_{i-1} < v^{\Delta}_{i-1} s_{\alpha_{t-i+1}}$.
- If $\Delta = \emptyset$, we have $v^{\Delta}_{i-1} = Id$, so that $v^{\Delta}_{i-1} < v^{\Delta}_{i-1} s_{\alpha_{t-i+1}}$.

□

Consider any $i \in [\![1, \dots ,t]\!]$, set $j = t - i + 1$ and $\{j_1 < \dots < j_s\} = \Delta \cap [\![j + 1, \dots ,t]\!] = \Delta \cap [\![t - i + 2, \dots ,t]\!]$, so that $v^{\Delta}_{i-1} = s^{\Delta}_{\alpha_t} \dots s^{\Delta}_{\alpha_{t-i+2}} = s_{\alpha_{j_s}} \dots s_{\alpha_{j_1}}$ and $v^{\Delta}_{i-1} s_{\alpha_{t-i+1}} = s_{\alpha_{j_s}} \dots s_{\alpha_{j_1}} s_{\alpha_j}$.

If w_1 and w_2 are in W, it results from ([12], corollary A.1.8) that

$(w_1 < w_2) \Leftrightarrow$ ($w_1 \neq w_2$ and w_1 is obtained by omitting some simple reflections in a reduced expression of w_2)

$\Leftrightarrow (w_1^{-1} < w_2^{-1})$.

So, with $w_1 = v^{\Delta}_{i-1}$ and $w_2 = v^{\Delta}_{i-1} s_{\alpha_{t-i+1}}$, we obtain:

$$(v^{\Delta}_{i-1} < v^{\Delta}_{i-1} s_{\alpha_{t-i+1}}) \quad \Leftrightarrow \quad (s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}} < s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}})$$

Let us set $u = s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}}$ and recall that $l(s_{\alpha_j} u) = l(u) \pm 1$. If $l(s_{\alpha_j} u) = l(u) + 1$, we have that $u < s_{\alpha_j} u$ ([12], A.1.6). Conversely, if $u < s_{\alpha_j} u$, we have $l(u) < l(s_{\alpha_j} u)$ ([12], A.1.6) and, consequently, $l(s_{\alpha_j} u) = l(u) + 1$. So, we have:

$$(v^{\Delta}_{i-1} < v^{\Delta}_{i-1} s_{\alpha_{t-i+1}}) \quad \Leftrightarrow \quad l(s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}}) = 1 + l(s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}})$$

Since the mapping $i \mapsto j = t - i + 1$ is a bijection of $[\![1, \dots, t]\!]$, we conclude:

Lemma 2.2. 4 *The diagram Δ is positive if and only if, for any $j \in [\![1, \dots, t]\!]$,*

$$\Delta \cap [\![j + 1, \dots, t]\!] = \{j_1 < \dots < j_s\} \Rightarrow l(s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}}) = 1 + l(s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}}).$$

Now, we can prove the following characterization of positive diagrams.

Proposition 2.2. 1 *The diagram Δ is positive if and only if, for any $j \in [\![1, \dots, t]\!]$,*

$$\Delta \cap [\![j + 1, \dots, t]\!] = \{j_1 < \dots < j_s\} \Rightarrow l(s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}}) = 1 + s.$$

Proof

Assume that Δ is positive, choose $j, j_1, \dots , j_s$ as above, and consider any l with $1 < l < s$. We have $\Delta \cap [\![j_l + 1, \dots , t]\!] = \{j_{l+1} < \dots < j_s\}$ and, by lemma 2.2. 4, $l(s_{\alpha_{j_l}} s_{\alpha_{j_{l+1}}} \dots s_{\alpha_{j_s}}) = 1 + l(s_{\alpha_{j_{l+1}}} \dots s_{\alpha_{j_s}})$. From this, we get that $l(s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}}) = s$ and, again by lemma 2.2. 4, $l(s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}}) = 1 + s$.
Conversely, assume that this equality holds for any j and choose $j, j_1, \dots , j_s$ as above. We have $\Delta \cap [\![j_1 + 1, \dots , t]\!] = \{j_2 < \dots < j_s\}$ and, by assumption, $l(s_{\alpha_{j_1}} s_{\alpha_{j_2}} \dots s_{\alpha_{j_s}}) = 1 + (s - 1) = s$. So, $l(s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}}) = 1 + l(s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}})$ and Δ is positive by lemma 2.2. 4.

□

2.3 Some properties of positive diagrams.

In this section, we use the same conventions as in section 2.2.

Proposition 2.3. 1

1. *The map:* $\Delta \mapsto u = v^{\Delta}$ *is a bijection from the set of positive diagrams onto the set* $\{u \in W \mid u \leq v\}$.
2. *The map:* $\Delta \mapsto u = w^{\Delta}$ *is a bijection from the set of positive diagrams onto the set* $\{u \in W \mid u \leq w\}$.

Proof

1. By lemma 2.2. 1, f: $\Delta \mapsto u = v^{\Delta}$ is a map from the set of positive diagrams in the set $\{u \in W \mid u \leq v\}$. Consider any $u \in W$ with $u \leq v$. By ([16], Lemma 3.5.), there exists a unique positive subexpression $\mathbf{u}^{+} = (u_0, u_1, \dots, u_t)$ of (6) with $u_t = u$. By lemma 2.2. 3, there exists a unique diagram Δ which satisfies $\mathbf{v}^{\Delta} = (v_0^{\Delta}, v_1^{\Delta}, \dots, v_t^{\Delta}) = \mathbf{u}^{+}$. So, Δ is positive (def. 2.2. 2) and $v^{\Delta} = v_t^{\Delta} = u_t = u$. This proves that f is surjective.
   If Δ' is a positive diagram such that $v^{\Delta'} = u$, then $\mathbf{v}^{\Delta'} = (v_0^{\Delta'}, v_1^{\Delta'}, \dots, v_t^{\Delta'})$ is a positive subexpression of (6) with $v_t^{\Delta'} = v^{\Delta'} = u$. So, we have $\mathbf{v}^{\Delta'} = \mathbf{u}^{+} \Rightarrow \mathbf{v}^{\Delta'} = \mathbf{v}^{\Delta}$ and, by lemma 2.2. 3, $\Delta' = \Delta$. This proves that f is bijective.

2. Denote by g the bijection $u \mapsto u^{-1}$ from $\{u \in W \mid u \leq w\}$ onto $\{u \in W \mid u \leq v\}$ (lemma 2.2. 1).
   Then $g^{-1} \circ f$: $\Delta \mapsto u = (v^{\Delta})^{-1} = w^{\Delta}$ is a bijection from the set of positive diagrams onto the set $\{u \in W \mid u \leq w\}$.

□

Consider any $p \in [\![1, \dots , t]\!]$, $p \neq 1, t$, and set

$$w_1 = s_{\alpha_1} \dots s_{\alpha_p} \tag{7}$$

$$w_2 = s_{\alpha_{p+1}} \dots s_{\alpha_t} \tag{8}$$

We have $w = w_1 w_2$ and, since (1) is a reduced expression of w, (7) and (8) are reduced expressions of w_1 and w_2 respectively.
Denote by Δ any subset of $[\![1, \dots , p]\!]$, so that Δ is a diagram with respect to (7) and with respect to (1) both.

Proposition 2.3. 2
(Δ *is positive with respect to* (7)) $\Leftrightarrow$ (Δ *is positive with respect to* (1))

Proof
Assume that Δ is positive with respect to (1) and consider any $j \in [\![1, \dots , p]\!]$. We have $\Delta \cap [\![j+1, \dots , p]\!] = \Delta \cap [\![j+1, \dots , t]\!]$ and, using the characterization of positive diagrams given in proposition 2.2. 1, we obtain that Δ is positive with respect to (7).
Assume that Δ is positive with respect to (7) and consider any $j \in [\![1, \dots , t]\!]$. If $j \leq p$, we have $\Delta \cap [\![j+1, \dots , t]\!] = \Delta \cap [\![j+1, \dots , p]\!]$, so that the characterization given in proposition 2.2. 1 is satisfied. If $j > p$, we have $\Delta \cap [\![j+1, \dots , t]\!] = \emptyset$, so that the characterization given in proposition 2.2. 1 is again satisfied.

□

Now, consider a non empty diagram Δ (with respect to (1)) and two integers j, m in $[\![1, \dots , t]\!]$ with $j < m$ and $m \in \Delta$. Let us recall that $(\beta_1, \dots , \beta_t)$ is the sequence of positive roots associated to the reduced expression (1)) of w (section 2.1) and let us denote:

- $\overline{\Delta} = [\![1, \dots , t]\!] \setminus \Delta$,
- $\Delta \cap [\![j+1, \dots , m-1]\!] = \{j_1 < \dots < j_r\}$ (unless this set is empty),
- $\overline{\Delta} \cap [\![j+1, \dots , m-1]\!] = \{l_1 < \dots < l_p\}$ (unless this set is empty),
- $(\gamma_1, \dots , \gamma_p, \gamma_{p+1})$ is the sequence of (non necessarily positive) roots defined recursively by $\gamma_{p+1} = \beta_m$ and, for $1 \leq i \leq p$, $\gamma_i = s_{\beta_{l_i}}(\gamma_{i+1})$,
- For $1 \leq i \leq p$, we set $a_i = (\beta_{l_i}^{\vee}, \gamma_{i+1})$.

Until the end of this section, we assume that $\overline{\Delta} \cap [\![j+1, \dots , m-1]\!]$ is nonempty, so that p and the roots γ_i $(1 \leq i \leq p+1)$ are well defined.

Lemma 2.3. 1

$$\gamma_1 = \beta_m - a_p\beta_{l_p} - \dots - a_1\beta_{l_1}$$

Proof
For $1 \leq i \leq p$, we have $\gamma_i = \gamma_{i+1} - a_i \beta_{l_i}$. Summing these equalities, we get $\gamma_1 = \gamma_{p+1} - a_p \beta_{l_p} - \dots - a_1 \beta_{l_1}$ and, by definition, $\gamma_{p+1} = \beta_m$.

$\square$

Set

$$w' = s_{\alpha_1} \dots s_{\alpha_{m-1}} \tag{9}$$

and observe that, since (1) is a reduced expression of w, (9) is a reduced expression of w'.

Lemma 2.3. 2
For any $i \in [\![1, \dots, p]\!]$, denote by w'_i the element of W obtained by omitting $s_{\alpha_{l_i}}, \dots, s_{\alpha_{l_p}}$ in (9).
Then, we have $\gamma_i = w'_i(\alpha_m)$.

Proof
Assume that $i = p$ and set $u_p = s_{\alpha_1} \dots s_{\alpha_{l_p - 1}}$, so that:
$w' = u_p s_{\alpha_{l_p}} s_{\alpha_{l_p+1}} \dots s_{\alpha_{m-1}} \Rightarrow u_p^{-1} w' = s_{\alpha_{l_p}} s_{\alpha_{l_p+1}} \dots s_{\alpha_{m-1}} \Rightarrow u_p s_{\alpha_{l_p}} u_p^{-1} w' = u_p s_{\alpha_{l_p+1}} \dots s_{\alpha_{m-1}} = w'_p$.
As $u_p(\alpha_{l_p}) = \beta_{l_p}$, we conclude that $w'_p = s_{\beta_{l_p}} w'$. So, $w'_p(\alpha_m) = s_{\beta_{l_p}} w'(\alpha_m) = s_{\beta_{l_p}}(\beta_m) = \gamma_p$.
Assume $i < p$, $\gamma_{i+1} = w'_{i+1}(\alpha_m)$, and set $u_i = s_{\alpha_1} \dots s_{\alpha_{l_i - 1}}$. We have $w'_{i+1} = u_i s_{\alpha_{l_i}} \sigma$ with $\sigma \in W$ and $w'_i = u_i \sigma$.
Form this, we deduce (as in the case $i = p$) that $u_i s_{\alpha_{l_i}} u_i^{-1} w'_{i+1} = w'_i$. So, $w'_i(\alpha_m) = s_{\beta_{l_i}} w'_{i+1}(\alpha_m) = s_{\beta_{l_i}}(\gamma_{i+1}) = \gamma_i$.

$\square$

Now, we can prove:

Proposition 2.3. 3 *Assume that*

$$\beta_j + \beta_m = a_p \beta_{l_p} + \dots + a_1 \beta_{l_1}.$$

Then Δ is not positive (with respect to (1)).

Proof
By lemma 2.3. 1, we have $\beta_j = -\gamma_1$ and, by lemma 2.3. 2, we can write $\beta_j = -w'_1(\alpha_m)$. Recall that w'_1 is obtained by omitting $s_{\alpha_{l_1}}, \dots, s_{\alpha_{l_p}}$ in the reduced expression (9) of w' and that
$\{l_1 < \dots < l_p\} = \overline{\Delta} \cap [\![j+1, \dots, m-1]\!] = [\![j+1, \dots, m-1]\!] \setminus \{j_1 < \dots < j_r\}$. So, we have

$$w'_1 = s_{\alpha_1} \dots s_{\alpha_{j-1}} s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_r}} = \eta s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_r}}$$

with $\eta = s_{\alpha_1} \dots s_{\alpha_{j-1}}$.
Since $\beta_j = \eta(\alpha_j)$, we have $\beta_j = -w'_1(\alpha_m) \Rightarrow \alpha_j = -s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_r}}(\alpha_m) \Rightarrow s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_r}}(\alpha_m) = -\alpha_j$ is a negative root. By ([9], lemma 10.2.C), this implies that $l(s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_r}} s_{\alpha_m}) < r + 2$.
As $m \in \Delta$, we have $\Delta \cap [\![j+1, \dots, t]\!] = \{j_1 < \dots < j_s\}$ with $s > r$ and $j_{r+1} = m$. So, $l(s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_s}}) \leq l(s_{\alpha_j} s_{\alpha_{j_1}} \dots s_{\alpha_{j_r}} s_{\alpha_m}) + s - (r+1) < r + 2 + s - (r+1) = s + 1$ and, by proposition 2.2. 1, Δ is not positive.

$\square$

3 Background on the deleting derivations algorithm.

3.1 Conventions.

In this section, we use the conventions of section 2.1 and we set $R = U^+[w]$. In order to simplify a little bit the notations, we set

- $X_i = X_{\beta_i}$ for any $i \in [\![1, \dots, t]\!]$, so that $R = k < X_1, \dots, X_t >$. Moreover, $X_1, \dots, X_t$ are called the canonical generators (with respect to the reduced decomposition (1)) of R.

- Recall (section 2.1) that, for each ρ in the root lattice $\mathbb{Z}\Pi$, the map h_ρ: $u \mapsto K_\rho.u$ is in $Aut(R)$, the group of automorphisms of the algebra R.

- Let us set $H = \{h_\rho \mid \rho \in \mathbb{Z}\Pi\}$ and observe that H is an abelian subgroup of $Aut(R)$.

- Recall that R is $\mathbb{Z}\Pi$-graded and that, for any homogeneous element u of degree γ in R, for any $\rho \in \mathbb{Z}\Pi$, we have $h_\rho(u) = q^{-(\rho,\gamma)}u$.
- For each $(i,j) \in (\llbracket 1, \dots, t \rrbracket)^2$, set $\lambda_{i,j} = q^{-(\beta_i,\beta_j)}$, $q_i = \lambda_{i,i} = q^{-\|\beta_i\|^2}$, and observe that q_i is not a root of unity.
- If $1 \leq i < j \leq t$, the Levendorskii-Soibeman formula can be written

$$X_jX_i - \lambda_{j,i}X_iX_j \ = \ P_{j,i}, \tag{10}$$

$$P_{j,i} \ = \sum_{\underline{a} \ = \ (a_{i+1},\dots,a_{j-1})} c_{\underline{a}}X_{i+1}^{a_{i+1}}\dots X_{j-1}^{a_{j-1}}. \tag{11}$$

  with $\underline{a} \in \mathbb{N}^{j-i-1}$, $c_{\underline{a}} \in k$, and $c_{\underline{a}} \neq 0$ for only finitely many $\underline{a}$. Moreover, $P_{j,i}$ is homogeneous with degree $\beta_i + \beta_j$ so that, if $j = i+1$, we have $P_{j,i} = 0$. In the general case, this also implies that, if $c_{\underline{a}} \neq 0$, then $a_{i+1}\beta_{i+1} \ + \ \dots \ + \ a_{j-1}\beta_{j-1} = \beta_i \ + \ \beta_j$.
- Since β_i and β_j are positive roots, $\beta_i \ + \ \beta_j$ is non zero. So, if $c_{\underline{a}} \neq 0$, then $\underline{a}$ is nonzero.
- From this, we get that the generators $X_1, \ \dots \ , \ X_t$ of R satisfy the equalities and the assumption 6.1.1. of ([4], section 6.1.).
- Moreover, since the ordered monomials $X^{\underline{a}} \ := X_1^{a_1} \ \dots \ X_t^{a_t}$, $\underline{a} = (a_1, \ \dots \ , a_1) \in \mathbb{N}^t$, are a basis of R, it results from ([4], propositions 6.1.1. and 6.1.2.) that R satisfies the conventions of ([4], section 3.1.). In particular, R is an iterated Ore extension of the ground field k and so, R is a noetherian domain.
- We denote by $F = Fract(R)$ the division ring of fractions of R.
- By ([4], proposition 6.1.1.), we also get that R is the k-algebra generated by the "variables" $X_1, \ \dots \ , \ X_t$ submitted to the relations (10).
- For each $l \in \llbracket 1, \dots, t \rrbracket$, we set $h_l = h_{\beta_l} \in H$, and observe that, if $i \in \llbracket 1, \dots, t \rrbracket$, we have $h_l(X_i) = \lambda_{l,i} \ X_i$.
- Since each X_i is homogeneous, it is an H-eigenvector and, since $\lambda_{1,1} = q^{-(\beta_1,\beta_1)}$ is not a root of unity, the assumption 4.1.2. of [4] is satisfied. Since each $\lambda_{i,j}$ is a power of q, the assumption 4.1.1. of [4] is also satisfied. As explained in [4] (proof of lemma 4.2.2.) this implies that each prime ideal of R is completely prime.
- Recall that each automorphism $h \in H$ can be extended in a (unique) automorphism (denoted h also) of F, so that H can be seen as a subgroup of $Aut(F)$.

3.2 The algebras $R^{(m)}$.

Recall ([4], section 3.) that, for any $m \in \llbracket 2, \dots, t+1 \rrbracket$, there exists a family $(X_1^{(m)}, \ \dots \ , \ X_t^{(m)})$ of new "variables" in F , called the canonical generators (with respect to the reduced decomposition (1)) of the algebra $R^{(m)} = k < X_1^{(m)}, \ \dots \ , \ X_t^{(m)} >$, and which satisfies the following properties:

- If $1 \leq i < j \leq t$, we have the following simplified Levendorskii-Soibelman formula:

$$X_j^{(m)}X_i^{(m)} - \lambda_{j,i}X_i^{(m)}X_j^{(m)} \ = \ P_{j,i}^{(m)} \tag{12}$$

with

$$\diamond \qquad m \leq j \ \Rightarrow \ P_{j,i}^{(m)} \ = \ 0. \tag{13}$$

$$\diamond \qquad j < m \ \Rightarrow \ P_{j,i}^{(m)} \ = \sum_{\underline{a} \ = \ (a_{i+1},\dots,a_{j-1})} c_{\underline{a}}(X_{i+1}^{(m)})^{a_{i+1}}\dots(X_{j-1}^{(m)})^{a_{j-1}} \tag{14}$$

where the coefficients $c_{\underline{a}}$ *are the same as in* (*11*).
(*So*, $P_{j,i}^{(m)} \ = \ 0$ *in the case* $i+1 = j < m$ and, in the general case, if $c_{\underline{a}} \neq 0$,
we have $a_{i+1}\beta_{i+1} + \dots + a_{j-1}\beta_{j-1} = \beta_i + \beta_i$, so that $\underline{a}$ is nonzero.)

- This implies that the generators $X_1^{(m)}, \dots, X_t^{(m)}$ of $R^{(m)}$ still satisfy the equalities and the assumption 6.1.1. of ([4], section 6.1.).

- The ordered monomials $(X^{(m)})^{\underline{a}} := (X_1^{(m)})^{a_1} \dots (X_t^{(m)})^{a_t}$, $\underline{a} = (a_1, \dots, a_1) \in \mathbb{N}^t$, are a basis of $R^{(m)}$. So, $R^{(m)}$ still satisfies the conventions of ([4], section 3.1.). In particular, $R^{(m)}$ is an iterated Ore extension of the ground field k and so, $R^{(m)}$ is a noetherian domain.

- We have $Fract(R^{(m)}) = Fract(R) = F$.

- By ([4], proposition 6.1.1.), $R^{(m)}$ is the k-algebra generated by the "variables" $X_1^{(m)}, \dots, X_t^{(m)}$ submitted to the relations (12).

- For each ρ in the root lattice $\mathbb{Z}\Pi$ and for each $i \in [\![1, \dots, t]\!]$, we still have $h_\rho(X_i^{(m)}) = q^{-(\rho,\beta_i)} X_i^{(m)}$. This implies that $h_\rho(R^{(m)}) = R^{(m)}$. So H can still be seen as a subgroup of $Aut(R^{(m)})$ and each $X_i^{(m)}$ is a H eigenvector. If l and i are in $[\![1, \dots, t]\!]$, we have $h_l(X_i^{(m)}) = \lambda_{l,i} X_i^{(m)}$ and, as above, this implies that $R^{(m)}$ satisfies the assumptions 4.4.2. and 4.4.1. of [4]. As explained in ([4], proof of lemma 4.2.2), this implies that each prime ideal of $R^{(m)}$ is completely prime.

- If $u \in F$, and if $\gamma \in \mathbb{Z}\Pi$, we say that u *is homogeneous of degree* γ, if $h_\rho(u) = q^{-(\rho,\gamma)}u$ for all ρ in $\mathbb{Z}\Pi$. So, for example, each $X_i^{(m)}$ is homogeneous of degree β_i.

- If $u_1, \dots, u_r$ are homogeneous of same degree γ, then any linear combination of $u_1, \dots, u_r$ (with coefficients in k) is homogeneous of degree γ.

- Clearly, if u_1 is homogeneous of degree γ_1 and u_2 is homogeneous of degree γ_2, then $u_1 u_2$ is homogeneous of degree $\gamma_1 + \gamma_2$.

- Likewise, if u is nonzero and homogeneous of degree γ, then u^{-1} is homogeneous of degree $-\gamma$.

- So, if $u_1, \dots, u_r$ are nonzero and homogeneous of degrees $\gamma_1, \dots, \gamma_r$ respectively, if $\underline{a} = (a_1, \dots, a_r) \in \mathbb{Z}^r$, then $u^{\underline{a}} := u_1^{a_1} \dots u_r^{a_r}$ is homogeneous of degrees $a_1\gamma_1 + \dots + a_r\gamma_r$.

- For each $j \in [\![1, \dots, t]\!]$, denote by $\delta_j^{(m)}$ the left h_j - derivation of $R^{(m)}$ defined by $X_j^{(m)}$. This means that, for each $a \in R^{(m)}$, $\delta_j^{(m)}(a) := X_j^{(m)}a - h_j(a)X_j^{(m)}$.

- If $j \in [\![1, \dots, t]\!]$, we have

$$h_j(k < X_1^{(m)}, \dots, X_{j-1}^{(m)} >) = k < X_1^{(m)}, \dots, X_{j-1}^{(m)} >,$$

$$\delta_j^{(m)}(k < X_1^{(m)}, \dots, X_{j-1}^{(m)} >) \subset k < X_1^{(m)}, \dots, X_{j-1}^{(m)} > .$$

So, h_j induces an automorphism, still denoted h_j, of the algebra $k < X_1^{(m)}, \dots, X_{j-1}^{(m)} >$. It is the only automorphism of $k < X_1^{(m)}, \dots, X_{j-1}^{(m)} >$ which satisfies $h_j(X_i^{(m)}) = \lambda_{j,i} X_i^{(m)}$ for any i.
Likewise, $\delta_j^{(m)}$ induces a left h_j - derivation of $k < X_1^{(m)}, \dots, X_{j-1}^{(m)} >$, still denoted $\delta_j^{(m)}$. It satisfies the following properties:

1. If $1 \leq i < j$, we have $\delta_j^{(m)}(X_i^{(m)}) = P_{j,i}^{(m)}$.
   This implies that $\delta_j^{(m)}$ is zero on $k < X_1^{(m)}, \dots, X_{j-1}^{(m)} >$ and so, that $X_j^{(m)}a = h_j(a)X_j^{(m)}$ for any a in $k < X_1^{(m)}, \dots, X_{j-1}^{(m)} >$, as soon as $j \geq m$.

2. $\delta_j^{(m)}$ is locally nilpotent on $k < X_1^{(m)}, \dots, X_{j-1}^{(m)} >$.

3. We have
$$R^{(m)} = k[X_1^{(m)}][X_2^{(m)}; h_2, \delta_2^{(m)}] \dots [X_{m-1}^{(m)}; h_{m-1}, \delta_{m-1}^{(m)}]\ [X_m^{(m)}; h_m] \dots [X_t^{(m)}; h_t].$$

- For each $i \in [\![1, \dots , t]\!]$, we have $X_i^{(t+1)} = X_i$, so that $R^{(t+1)} = R$.

Until the end of this section, we assume that $m \in [\![2, \dots , t]\!]$.

- Recall that $q_m = \lambda_{m,m}$ is not a root of unity and define quantum integers $[l]_{q_m}$ (resp. quantum factorial $[l]!_{q_m}$) for $l \in \mathbb{N}$, as in ([4], section 2.).
- For each $i \in [\![1, \dots , t]\!]$, we have
  1. $m \leq i \Rightarrow X_i^{(m)} = X_i^{(m+1)}$. In particular, we have $X_m^{(m)} = X_m^{(m+1)}$.
  2. $$i < m \Rightarrow X_i^{(m)} = X_i^{(m+1)} + \sum_{l=1}^{+\infty} C_l^{(m+1)} (X_m^{(m+1)})^{-l}$$
     with
     $$C_l^{(m+1)} = \frac{(1-q_m)^{-l}}{[l]!_{q_m}} \lambda_{m,i}^{-l} (\delta_m^{(m+1)})^l (X_i^{(m+1)}).$$
- We observe that, since $k < X_1^{(m+1)}, \dots , X_{m-1}^{(m+1)} >$ is $\delta_m^{(m+1)}$ - stable, we have $C_l^{(m+1)} \in k < X_1^{(m+1)}, \dots , X_{m-1}^{(m+1)} >$ and, since $\delta_m^{(m+1)}$ is locally nilpotent on $k < X_1^{(m+1)}, \dots , X_{m-1}^{(m+1)} >$, only finitely many $C_l^{(m+1)}$ are non zero.
- If $i < m$ and $X_m^{(m+1)} X_i^{(m+1)} - \lambda_{m,j} X_i^{(m+1)} X_m^{(m+1)} = P_{m,i}^{(m+1)} = 0$, then $\delta_m^{(m+1)}(X_i^{(m+1)}) = 0$ and so, $X_i^{(m)} = X_i^{(m+1)}$.
- There exists a unique homomorphism of k-algebras
  $$\Theta^{(m)} : k < X_1^{(m+1)}, \dots , X_{m-1}^{(m+1)} > \rightarrow k < X_1^{(m)}, \dots , X_{m-1}^{(m)} >$$
  which verifies $\Theta^{(m)}(X_i^{(m+1)}) = X_i^{(m)}$ for $1 \leq i \leq m-1$.
- When $m = 2$, we set
  1. $Z_i = X_i^{(2)}$ for $i \in [\![1, \dots , t]\!]$.
  2. $\overline{R} = R^{(2)}$.
- $\overline{R} = k < Z_1, \dots , Z_t >$ is the k - algebra generated by the variables $Z_1, \dots , Z_t$ submitted to the relations:
  $$(1 \leq i < j \leq t) \Rightarrow Z_j Z_i = \lambda_{j,i} Z_i Z_j \tag{15}$$
- For any $i \in [\![1, \dots , t]\!]$, we have $X_i^{(2)} = \dots = X_i^{(i+1)} = Z_i$.
- $S_m = \{(X_m^{(m+1)})^l \mid l \in \mathbb{N}\} = \{(X_m^{(m)})^l \mid l \in \mathbb{N}\}$ is a multiplicative system of regular elements in $R^{(m)}$ and in $R^{(m+1)}$. It satisfies the Ore condition (on both sides) in each one of those rings and we have $R^{(m)} S_m^{-1} = R^{(m+1)} S_m^{-1}$.

3.3 Prime and H-prime spectrum of $R^{(m)}$, admissible diagrams

Consider any integer $m \in [\![2, \dots , t+1]\!]$. As usual, we denote by $Spec(R^{(m)})$ (resp. $H - Spec(R^{(m)})$) the set of all prime ideals (resp. H- invariant prime ideals) of $R^{(m)}$. Recall (see section 3.2) that each prime ideal of $R^{(m)}$ is completely prime. If $m \in [\![2, \dots , t]\!]$, we denote by

$$\phi_m : Spec(R^{(m+1)}) \rightarrow Spec(R^{(m)})$$

the canonical injection defined in ([4], section 4.3).
Moreover, if $m' \leq m+1$ and if $\mathcal{P} \in Spec(R^{(m+1)})$, then $\mathcal{P}' = \phi_{m'} \circ \dots \circ \phi_m(\mathcal{P})$ is called the canonical image of $\mathcal{P}$ in $Spec(R^{(m')})$. (If $m' = m+1$, this canonical image is $\mathcal{P}$ itself.)

Let us now recall the main properties of the maps ϕ_m.

- Denote by $\mathcal{P}^{(m+1)}$ any prime ideal in $Spec(R^{(m+1)})$ and set $\mathcal{P}^{(m)} = \phi_m(\mathcal{P}^{(m+1)})$.
  1. If $X_m^{(m+1)} \notin \mathcal{P}^{(m+1)}$, then $\mathcal{P}^{(m+1)} \cap S_m = \emptyset$ and
  $$\mathcal{P}^{(m)} = R^{(m)} \cap \mathcal{P}^{(m+1)} S_m^{-1}.$$
  In this case, $X_m^{(m)} = X_m^{(m+1)} \notin \mathcal{P}^{(m)}$.
  2. If $X_m^{(m+1)} \in \mathcal{P}^{(m+1)}$, there exists a unique algebra homomorphism $g : R^{(m)} \to R^{(m+1)}/\mathcal{P}^{(m+1)}$ which verifies $g(X_i^{(m)}) = X_i^{(m+1)} + \mathcal{P}^{(m+1)}$ for all $i \in [\![1, \dots , t]\!]$, and $\mathcal{P}^{(m)} = Ker(g)$.
  In this case, $X_m^{(m)} = X_m^{(m+1)} \in \mathcal{P}^{(m)}$.
  3. For any $h \in H$, we have $h(\mathcal{P}^{(m)}) = \phi_m(h(\mathcal{P}^{(m+1)}))$ ([4], lemma 5.5.5.) so that, by the injectivity of ϕ_m, $\mathcal{P}^{(m+1)}$ is H - invariant if and only if $\mathcal{P}^{(m)}$ is H - invariant.
- A diagram Δ is said admissible (with respect to the reduced decomposition (1)) if there exists a prime ideal $\mathcal{P}$ of R ($= R^{(t+1)}$) whose canonical image $\mathcal{P}^{(2)}$ in $Spec(\overline{R})$ ($= Spec(R^{(2)})$) verifies:
$$\mathcal{P}^{(2)} \cap \{Z_i \mid 1 \leq i \leq t\} = \{Z_i \mid i \in \Delta\}.$$
- Consider a diagram Δ and denote by $\mathcal{P}_\Delta^{(2)} = (\{Z_i \mid i \in \Delta\})$ the ideal generated by the variables Z_i with $i \in \Delta$. Then, by ([4], proposition 5.5.1.), we have
  1. $\mathcal{P}_\Delta^{(2)} \in H - Spec(R^{(2)})$.
  2. Conversely, for any $\mathcal{Q} \in H - Spec(R^{(2)})$, there exists a (unique) diagram Δ such that $\mathcal{Q} = \mathcal{P}_\Delta^{(2)}$, namely $\Delta = \{i \in [\![1, t]\!] \mid Z_i \in Q\}$.
- A diagram Δ is admissible if and only if there exists $\mathcal{P}_\Delta \in Spec(R)$ $(= Spec(R^{(t+1)}))$ such that
$$\mathcal{P}_\Delta^{(2)} = \phi_2 \circ \dots \circ \phi_t(\mathcal{P}_\Delta).$$
(See [4], theorem 5.5.1. and observe that, since each ϕ_i is injective, $\mathcal{P}_\Delta$ is unique.)
- The map $\Delta \mapsto \mathcal{P}_\Delta$ is a bijection from the set of admissible diagrams onto the set $H - Spec(R)$ $(= H - Spec(R^{(t+1)}))$. In fact, if Δ is admissible, then $\mathcal{P}_\Delta$ is H - invariant because $\mathcal{P}_\Delta^{(2)} = \phi_2 \circ \dots \circ \phi_t(\mathcal{P}_\Delta)$ is. So, $\Delta \mapsto \mathcal{P}_\Delta$ is a map from the set of admissible diagrams into $H - Spec(R)$. It is injective because the map $\Delta \mapsto \mathcal{P}_\Delta^{(2)}$ is injective. If $\mathcal{P} \in H - Spec(R)$, then $\phi_2 \circ \dots \circ \phi_t(\mathcal{P}) \in H - Spec(R^{(2)})$. So $\phi_2 \circ \dots \circ \phi_t(\mathcal{P}) = \mathcal{P}_\Delta^{(2)}$ with Δ an admissible diagram such that $\mathcal{P} = \mathcal{P}_\Delta$.

4 New results on H-invariant prime ideals.

In this section, we consider an integer $m \in [\![2, \dots , t+1]\!]$, and we denote by $\mathcal{P}^{(m)}$ an H-invariant prime ideal. We set $A^{(m)} = R^{(m)}/P^{(m)}$ and we observe that this algebra is a noetherian domain (since $R^{(m)}$ is noetherian and $P^{(m)}$ is completely prime). Set $D_m = Frac(A^{(m)})$ it's division ring of fractions. Denote by $f_m : R^{(m)} \to A^{(m)}$ the canonical homomorphism and, for each $i \in [\![1, \dots , t]\!]$, set $x_i^{(m)} = f_m(X_i^{(m)})$ the canonical image of $X_i^{(m)}$ in $A^{(m)}$. If h is an element of the group H, we have $h(\mathcal{P}^{(m)}) = \mathcal{P}^{(m)}$ and, consequently, h induces an automorphism of the algebra $A^{(m)}$, denoted $\overline{h}$, which satisfies $\overline{h} \circ f_m = f_m \circ h$. This automorphism can be extended in an automorphism, still denoted $\overline{h}$ of the division ring D_m.

4.1 A necessary and sufficient condition for $\mathcal{P}^{(m)}$ to be in $Im(\phi_m)$.

Assume that $m \leq t$ and recall (section 3.2) that $R^{(m)}$ is the k - algebra generated by the variables $X_1^{(m)}, \dots , X_t^{(m)}$ submitted to the relations 12:
$$X_j^{(m)} X_i^{(m)} - \lambda_{j,i} X_i^{(m)} X_j^{(m)} = P_{j,i}^{(m)}(X_{i+1}^{(m)}, \dots , X_{j-1}^{(m)})$$
for $i < j$, and with $P_{j,i}^{(m)}(X_{i+1}^{(m)}, \dots , X_{j-1}^{(m)}) \in k < X_{i+1}^{(m)}, \dots , X_{j-1}^{(m)} >$.

Proposition 4.1. 1
$\mathcal{P}^{(m)} \in Im(\phi_m)$ if and only if one of two following conditions is satisfied
a) $X_m^{(m)} \notin \mathcal{P}^{(m)}$.
b) $X_m^{(m)} \in \mathcal{P}^{(m)}$ and $\Theta^{(m)}(\delta_m^{(m+1)}(X_i^{(m+1)})) \in \mathcal{P}^{(m)}$ for $1 \leq i \leq m-1$.

Proof.

Assume that $\mathcal{P}^{(m)} \in Im(\phi_m)$, so that $\mathcal{P}^{(m)} = \phi_m(\mathcal{P}^{(m+1)})$ with $\mathcal{P}^{(m+1)} \in Spec(R^{(m+1)})$, and assume that condition a) is not satisfied. This implies that $\mathcal{P}^{(m)} = ker(g)$ where $g : R^{(m)} \rightarrow R^{(m+1)}/\mathcal{P}^{(m+1)}$ is the homomorphism which transforms each $X_i^{(m)}$ in $x_i^{(m+1)} = X_i^{(m+1)} + \mathcal{P}^{(m+1)}$. Consider $1 \leq i \leq m-1$.
Recall that $\delta_m^{(m+1)}(X_i^{(m+1)}) = P_{m,i}^{(m+1)}(X_{i+1}^{(m+1)}, \ldots, X_{m-1}^{(m+1)})$ and and that
$\Theta^{(m)} : k < X_1^{(m+1)}, \ldots, X_{m-1}^{(m+1)} > \rightarrow k < X_1^{(m)}, \ldots, X_{m-1}^{(m)} >$ is the homomorphism which transforms each $X_l^{(m+1)}$ in $X_l^{(m)}$. Since $X_m^{(m)} \in \mathcal{P}^{(m)}$, we have $X_m^{(m+1)} \in \mathcal{P}^{(m+1)}$ ([4], proposition 4.3.1.) and so, $\delta_m^{(m+1)}(X_i^{(m+1)}) \in \mathcal{P}^{(m+1)}$. Now, we have
$g(\Theta^{(m)}(\delta_m^{(m+1)}(X_i^{(m+1)}))) = g(\Theta^{(m)}(P_{m,i}^{(m+1)}(X_{i+1}^{(m+1)}, \ldots, X_{m-1}^{(m+1)}))) = g(P_{m,i}^{(m+1)}(X_{i+1}^{(m)}, \ldots, X_{m-1}^{(m)})) = P_{m,i}^{(m+1)}(x_{i+1}^{(m+1)}, \ldots, x_{m-1}^{(m+1)}) = P_{m,i}^{(m+1)}(X_{i+1}^{(m+1)}, \ldots, X_{m-1}^{(m+1)}) + \mathcal{P}^{(m+1)} = 0$.
This implies that $\Theta^{(m)}(\delta_m^{(m+1)}(X_i^{(m+1)}))) \in ker(g) = \mathcal{P}^{(m)}$.

If condition a) is satisfied, then $\mathcal{P}^{(m)} \in Im(\phi_m)$ by ([4], lemma 4.3.1.).

Assume that condition b) is satisfied. So, if $1 \leq i \leq m-1$, we have, as previously, $P_{m,i}^{(m+1)}(X_{i+1}^{(m)}, \ldots, X_{m-1}^{(m)}) = \Theta^{(m)}(\delta_m^{(m+1)}(X_i^{(m+1)})) \in \mathcal{P}^{(m)}$. So, in $A^{(m)} = R^{(m)}/\mathcal{P}^{(m)}$, we have $P_{m,i}^{(m+1)}(x_{i+1}^{(m)}, \ldots, x_{m-1}^{(m)}) = 0$.
Since $P_{m,i}^{(m)} = 0$ (see section 3.2), we can write $x_m^{(m)}x_i^{(m)} - \lambda_{m,i}x_i^{(m)}x_m^{(m)} = P_{m,i}^{(m)}(x_{i+1}^{(m)}, \ldots, x_{m-1}^{(m)}) = 0 = P_{m,i}^{(m+1)}(x_{i+1}^{(m)}, \ldots, x_{m-1}^{(m)})$.
If $1 \leq i \leq j-1$ with $j \neq m$, we have (see section 3.2)
$x_j^{(m)}x_i^{(m)} - \lambda_{j,i}x_i^{(m)}x_j^{(m)} = P_{j,i}^{(m)}(x_{i+1}^{(m)}, \ldots, x_{j-1}^{(m)}) = P_{j,i}^{(m+1)}(x_{i+1}^{(m)}, \ldots, x_{j-1}^{(m)})$.
So, by the universal property of algebras defined by generators and relations, there exists a (unique) homomorphism $\epsilon : R^{(m+1)} \rightarrow A^{(m)}$ which transforms each $X_l^{(m+1)}$ in $x_l^{(m)}$. This homomorphism is surjective, and its kernel $ker(\epsilon) = \mathcal{P}^{(m+1)}$ is a prime ideal of $R^{(m+1)}$. We observe that, since $X_m^{(m)} \in \mathcal{P}^{(m)}$, we have $X_m^{(m+1)} \in \mathcal{P}^{(m+1)}$, and that ϵ induces an automorphism

$$\overline{\epsilon} : R^{(m+1)}/\mathcal{P}^{(m+1)} \rightarrow A^{(m)} = R^{(m)}/\mathcal{P}^{(m)}$$

which transforms each $x_l^{(m+1)}$ in $x_l^{(m)}$. Recall that $f_m : R^{(m)} \rightarrow R^{(m)}/\mathcal{P}^{(m)}$ denotes the canonical homomorphism. So, $g = (\overline{\epsilon})^{-1} \circ f_m : R^{(m)} \rightarrow R^{(m+1)}/\mathcal{P}^{(m+1)}$ is the homomorphism which transforms each $X_l^{(m)}$ in $x_l^{(m+1)}$. As $ker(g) = ker(f_m) = \mathcal{P}^{(m)}$, we conclude that $\mathcal{P}^{(m)} = \phi_m(\mathcal{P}^{(m+1)})$.

□

Corollary 4.1. 1
Assume $2 \leq m \leq t+1$ and consider some $\mathcal{Q} \in Spec(R^{(m)})$.
Assume that $X_m^{(m)} \notin \mathcal{Q}, \ldots, X_t^{(m)} \notin \mathcal{Q}$.
Then there exists $\mathcal{P} \in Spec(R)$ such that $\mathcal{Q}$ is the canonical image of $\mathcal{P}$ in $Spec(R^{(m)})$. Moreover, if $\mathcal{P}^{(2)}$ is the canonical image of $\mathcal{P}$ in $Spec(R^{(2)})$, then

$$\mathcal{P}^{(2)} \cap \{Z_m, \ldots, Z_t\} = \emptyset.$$

Proof.
We prove this by decreasing induction on m.

- If $m = t+1$, we have $\mathcal{Q} \in Spec(R)$ and $\mathcal{P} = \mathcal{Q}$ satisfies the required properties.

- Assume $2 \leq m \leq t$. By proposition 4.1. 1, we have $\mathcal{Q} = \phi_m(\mathcal{Q}')$ with $\mathcal{Q}' \in Spec(R^{(m+1)})$. Since $X_m^{(m)} \notin \mathcal{Q}$, we have (see section 3.3) $X_m^{(m+1)} \notin \mathcal{Q}'$ and, if we set $S_m = \{(X_m^{(m+1)})^d \mid d \in \mathbb{N}\}$,

$$\mathcal{Q} \ = \ R^{(m)} \ \cap \ \mathcal{Q}' S_m^{-1}.$$

Assume $X_j^{(m+1)} \in \mathcal{Q}'$ with $j \geq m$. Since $X_j^{(m+1)} = X_j^{(m)}$ (see section 3.2), we have $X_j^{(m)} \in R^{(m)} \ \cap \ \mathcal{Q}' \subseteq R^{(m)} \ \cap \ \mathcal{Q}' S_m^{-1} = \mathcal{Q}$, which is false. So, we have:

- $X_m^{(m+1)} \notin \mathcal{Q}'$.
- $X_{m+1}^{(m+1)} \notin \mathcal{Q}', \ \dots \ , \ X_t^{(m+1)} \notin \mathcal{Q}'$ and, by the induction assumption, there exists $\mathcal{P} \in Spec(R)$ such that $\mathcal{Q}'$ is the canonical image of $\mathcal{P}$ in $Spec(R^{(m+1)})$. This implies that $\mathcal{Q} = \phi_m(\mathcal{Q}')$ is the canonical image of $\mathcal{P}$ in $Spec(R^{(m)})$. Moreover, again by the induction assumption, we have

$$\mathcal{P}^{(2)} \ \cap \ \{Z_{m+1}, \ \dots \ , \ Z_t\} \ = \ \emptyset.$$

As $X_m^{(m+1)} \notin \mathcal{Q}'$, we know ([4], proposition 5.2.1.) that $Z_m \notin \mathcal{P}^{(2)}$.

□

4.2 Some properties of $A^{(m)}$.

For each integer $j \in [\![2, \dots , m-1]\!]$, we set $\mathcal{P}^{(j)} \ = \ \phi_j \circ \ \dots \ \circ \phi_{m-1}(\mathcal{P}^{(m)})$. Let us, first, construct in $A^{(m)}$, a new version of the deleting derivations algorithm.

Proposition 4.2. 1

For each integer $j \in [\![2, \dots , m-1]\!]$, there exists a sequence $(x_1^{(j)}, \ \dots \ , \ x_t^{(j)})$ in $D_m \ = \ Frac(A^{(m)})$ and a subalgebra $A^{(j)} \ = k < x_1^{(j)}, \ \dots \ , \ x_t^{(j)} >$ of D_m which satisfy the following properties.

1. *If l and i are in $[\![1, \dots , t]\!]$, we have $\overline{h_l}(x_i^{(j)}) \ = \ \lambda_{l,i} x_i^{(j)}$, $\overline{h_l}(A^{(j)}) \ = \ A^{(j)}$ and $\overline{h_l}$ induces, by restriction, an automorphism (still denoted $\overline{h_l}$) of $A^{(j)}$.*
2. *Choose $l \in [\![1, \dots , t]\!]$ and denote by $d_l^{(j)}$ the left $\overline{h_l}$ - derivation of $A^{(j)}$ associated to $x_l^{(j)}$ ($d_l^{(j)}(a) \ = \ x_l^{(j)} a - \overline{h_l}(a) x_l^{(j)}$ for all $a \ \in \ A^{(j)}$). Then $d_l^{(j)}(k < x_1^{(j)}, \ \dots \ , \ x_{l-1}^{(j)} >) \ \subset \ k < x_1^{(j)}, \ \dots \ , \ x_{l-1}^{(j)} >$ and $d_l^{(j)}$ is locally nilpotent on $k < x_1^{(j)}, \ \dots \ , \ x_{l-1}^{(j)} >$.*
3. *There exists a unique homomorphism $f_j : \ R^{(j)} \ \rightarrow \ A^{(j)}$ which transforms each $X_i^{(j)}$ in $x_i^{(j)}$. f_j is surjective and $ker(f_j) \ = \ \mathcal{P}^{(j)}$.*
4. *If $l \in [\![1, \dots , t]\!]$, we have $d_l^{(j)} \circ f_j \ = \ f_j \circ \delta_l^{(j)}$ and $\overline{h_l} \circ f_j \ = \ f_j \circ h_l$.*
5. *If $l, e \in [\![1, \dots , t]\!]$, we have $\overline{h_l} \circ \overline{h_e} \ = \ \overline{h_e} \circ \overline{h_l}$ and $\overline{h_l} \circ d_e^{(j)} \ = \ \lambda_{l,e} d_e^{(j)} \circ \overline{h_l}$.*
6. *If $x_j^{(j+1)} \ = \ 0$, then $x_i^{(j)} \ = \ x_i^{(j+1)}$ for each $i \in [\![1, \dots , t]\!]$.*
7. *If $x_j^{(j+1)} \ \neq \ 0$, and $i \in [\![1, \dots , t]\!]$, then*
    (a) $j \leq i \ \Rightarrow \ x_i^{(j)} \ = \ x_i^{(j+1)}$. In particular, we have $x_j^{(j)} \ = \ x_j^{(j+1)}$.
    (b) $i < j \ \Rightarrow \ x_i^{(j)} \ = \ x_i^{(j+1)} \ + \ \sum_{d=1}^{+\infty} c_d^{(j+1)} (x_j^{(j+1)})^{-d}$ with $c_d^{(j+1)} \ = \ \dfrac{(1-q_j)^{-d}}{[d]!_{q_j}} \ \lambda_{j,i}^{-d} \ (d_j^{(j+1)})^d \ (x_i^{(j+1)})$.

    (We observe that, since $k < x_1^{(j+1)}, \ \dots \ , \ x_{j-1}^{(j+1)} >$ is $d_j^{(j+1)}$ - stable, we have $c_d^{(j+1)} \in \ k < x_1^{(j+1)}, \ \dots \ , \ x_{j-1}^{(j+1)} >$ and, since $d_j^{(j+1)}$ is locally nilpotent on $k < x_1^{(j+1)}, \ \dots \ , \ x_{j-1}^{(j+1)} >$, only finitely many $c_d^{(j+1)}$ are non zero.)

Proof.

First observe that, if 3. is satisfied and if $\overline{h_l} \circ f_j \ = \ f_j \circ h_l$ (as homomorphism from $R^{(j)}$ towards $D^{(m)}$) for any l, then properties 1. to 5. are all satisfied. In fact,

1. For all l and i in $[\![1, \dots, t]\!]$, we have $\overline{h_l}(x_i^{(j)}) = \overline{h_l}\circ f_j(X_i^{(j)}) = f_j\circ h_l(X_i^{(j)}) = \lambda_{l,i}f_j(X_i^{(j)}) = \lambda_{l,i}x_i^{(j)}$. This immediately implies that $\overline{h_l}(A^{(j)}) = A^{(j)}$ and $\overline{h_l}$ induces, by restriction, an automorphism of $A^{(j)}$.

4. If $P \in R^{(j)}$, we have $d_l^{(j)}(f_j(P)) = x_l^{(j)}f_j(P) - \overline{h_l}(f_j(P))x_l^{(j)}, = f_j(X_l^{(j)}P - h_l(P)X_l^{(j)}) = f_j\circ\delta_l^{(j)}(P)$. This proves that $d_l^{(j)}\circ f_j = f_j\circ\delta_l^{(j)}$.

2. By 4., we have $d_l^{(j)}(k < x_1^{(j)}, \dots, x_{l-1}^{(j)} >) = d_l^{(j)}(f_j(k < X_1^{(j)}, \dots, X_{l-1}^{(j)} >)) = f_j(\delta_l^{(j)}(k < X_1^{(j)}, \dots, X_{l-1}^{(j)} >)) \subset f_j(k < X_1^{(j)}, \dots, X_{l-1}^{(j)} >) = k < x_1^{(j)}, \dots, x_{l-1}^{(j)} >$. Moreover, if $a \in k < x_1^{(j)}, \dots, x_{l-1}^{(j)} >$, we can write $a = f_j(P)$ with $P \in k < X_1^{(j)}, \dots, X_{l-1}^{(j)} >$. Since $\delta_l^{(j)}$ is locally nilpotent on $k < X_1^{(j)}, \dots, X_{l-1}^{(j)} >$, we have $(\delta_l^{(j)})^r(P) = 0$ for some $r \in \mathbb{N}$, and $(d_l^{(j)})^r(a) = (d_l^{(j)})^r\circ f_j(P) = f_j\circ(\delta_l^{(j)})^r(P) = 0$. So, $d_l^{(j)}$ is locally nilpotent on $k < x_1^{(j)}, \dots, x_{l-1}^{(j)} >$.

5. If i in $[\![1, \dots, t]\!]$, we have $\overline{h_l}\circ\overline{h_e}(x_i^{(j)}) = \lambda_{l,i}\lambda_{e,i}x_i^{(j)} = \overline{h_e}\circ\overline{h_l}(x_i^{(j)})$ and, since $A^{(j)}$ is generated by the elements $x_i^{(j)}$, we have $\overline{h_l}\circ\overline{h_e} = \overline{h_e}\circ\overline{h_l}$.
Consider two elements a, b in $A^{(j)}$ such that the maps $\overline{h_l}\circ d_e^{(j)}$ and $\lambda_{l,e}d_e^{(j)}\circ\overline{h_l}$ coincide on a and b. Then, we have $\overline{h_l}\circ d_e^{(j)}(ab) = \overline{h_l}(d_e^{(j)}(a)b + \overline{h_e}(a)d_e^{(j)}(b)) = \lambda_{l,e}d_e^{(j)}(\overline{h_l}(a))\overline{h_l}(b) + \lambda_{l,e}\overline{h_e}(\overline{h_l}(a))d_e^{(j)}(\overline{h_l}(b)) = \lambda_{l,e}d_e^{(j)}\circ\overline{h_l}(ab)$. So, we just have to prove that $\overline{h_l}\circ d_e^{(j)}$ and $\lambda_{l,e}d_e^{(j)}\circ\overline{h_l}$ coincide on each $x_i^{(j)}$. Now, $\overline{h_l}\circ d_e^{(j)}(x_i^{(j)}) = \overline{h_l}(x_e^{(j)}x_i^{(j)} - \lambda_{e,i}x_i^{(j)}x_e^{(j)}) = \lambda_{l,e}\lambda_{l,i}(x_e^{(j)}x_i^{(j)} - \lambda_{e,i}x_i^{(j)}x_j^{(j)}) = \lambda_{l,e}\lambda_{l,i}d_e^{(j)}(x_i^{(j)}) = \lambda_{l,e}d_e^{(j)}(\overline{h_l}(x_i^{(j)}))$. So we can conclude that $\overline{h_l}\circ d_e^{(j)} = \lambda_{l,e}d_e^{(j)}\circ\overline{h_l}$.

For $j \in [\![2, \dots, m-1]\!]$, denote by $\mathcal{H}_j$ the following assumption:

There exists a sequence $(x_1^{(j+1)}, \dots, x_t^{(j+1)}) \in (D_m)^t$ *and a subalgebra* $A^{(j+1)} = k < x_1^{(j+1)}, \dots, x_t^{(j+1)} >$ *of* D_m *which satisfy the analog of properties 1. to 5. obtained by changing* j *in* $j+1$.

Observe that the assumption $\mathcal{H}_{m-1}$ is satisfied. In fact, the constructions of f_m and of the variables $x_i^{(m)}$ (see the beginning of section 4) imply straightway the property 3. and the equality $\overline{h}\circ f_m = f_m\circ h$ for any h in H.
Now, assume that $\mathcal{H}_j$ is satisfied, define the elements $x_i^{(j)}$ as in 6. and 7. and set $A^{(j)} = k < x_1^{(j)}, \dots, x_t^{(j)} >$. It remains to prove that property 3. is satisfied and that $\overline{h_l}\circ f_j = f_j\circ h_l$.

- Assume that $x_j^{(j+1)} = 0$, so that $X_j^{(j+1)} \in Ker(f_{j+1}) = \mathcal{P}^{(j+1)}$. Recall that, in this case, $\mathcal{P}^{(j)}$ is the kernel of the homomorphism $g: R^{(j)} \to R^{(j+1)}/\mathcal{P}^{(j+1)}$ which transforms each $X_i^{(j)}$ in $X_i^{(j+1)} + \mathcal{P}^{(j+1)}$.
By the assumption $\mathcal{H}_j$, $f_{j+1}: R^{(j+1)} \to A^{(j+1)}$ induces an isomorphism $\widehat{f_{j+1}}: R^{(j+1)}/\mathcal{P}^{(j+1)} \to A^{(j+1)}$ which transforms each $X_i^{(j+1)} + \mathcal{P}^{(j+1)}$ in $x_i^{(j+1)}$. So $f_j = \widehat{f_{j+1}}\circ g: R^{(j)} \to A^{(j+1)}$ is the homomorphism which transforms each $X_i^{(j)}$ in $x_i^{(j+1)}$, and we have $ker(f_j) = ker(g) = \mathcal{P}^{(j)}$. Since, in this case, we have $x_i^{(j)} = x_i^{(j+1)}$ for each i, we obtain that f_j satisfies the property 3. Moreover, if l and i are in $[\![1, \dots, t]\!]$, we have $\overline{h_l}\circ f_j(X_i^{(j)}) = \overline{h_l}(x_i^{(j)}) = \overline{h_l}(x_i^{(j+1)}) = \lambda_{l,i}(x_i^{(j+1)}) = \lambda_{l,i}(x_i^{(j)}) = f_j\circ h_l(X_i^{(j)})$. So, $\overline{h_l}\circ f_j = f_j\circ h_l$.
- Assume that $x_j^{(j+1)} \neq 0$, so that $X_j^{(j+1)} \notin Ker(f_{j+1}) = \mathcal{P}^{(j+1)}$. Set $S_j = \{(X_j^{(j+1)})^l \mid l \in \mathbb{N}\}$ and recall that
$$\mathcal{P}^{(j)} = \mathcal{P}^{(j+1)}S_j^{-1}\cap R^j.$$
As $f_{j+1}(X_j^{(j+1)}) = x_j^{(j+1)}$ is invertible in D_m, f_{j+1} can be extended in an homomorphism $\widetilde{f_{j+1}}: R^{(j+1)}S_j^{-1} \to D_m$ $(\widetilde{f_{j+1}}(Q(X_j^{(j+1)})^{-l}) = f_{j+1}(Q)(x_j^{(j+1)})^{-l}$ for all Q in $R^{(j+1)}$ and l in $\mathbb{N})$ and we have $ker(\widetilde{f_{j+1}}) = \mathcal{P}^{(j+1)}S_j^{-1}$. Recall that $R^{(j)} \subset R^{(j+1)}S_j^{-1}$ and let us compute $\widetilde{f_{j+1}}(X_i^{(j)})$ for each $i \in [\![1, \dots, t]\!]$.

  - If $j \leq i$, then we have $X_i^{(j)} = X_i^{(j+1)}$ (see section 3.2) and $\widetilde{f_{j+1}}(X_i^{(j)}) = x_i^{(j+1)} = x_i^{(j)}$.
  - If $i < j$, then we have
$$X_i^{(j)} = X_i^{(j+1)} + \sum_{d=1}^{+\infty} C_d^{(j+1)}(X_j^{(j+1)})^{-d}$$

with

$$C_d^{(j+1)} = \frac{(1-q_j)^{-d}}{[d]!_{q_j}} \lambda_{j,i}^{-d} \, (\delta_j^{(j+1)})^d \, (X_i^{(j+1)}).$$

As $C_d^{(j+1)} \in R^{(j+1)}$, we have $\widetilde{f_{j+1}}(C_d^{(j)}) = f_{j+1}(C_d^{(j)}) = \frac{(1-q_j)^{-d}}{[d]!_{q_j}} \lambda_{j,i}^{-d} \, f_{j+1} \circ (\delta_j^{(j+1)})^d \, (X_i^{(j+1)}) = \frac{(1-q_j)^{-d}}{[d]!_{q_j}} \lambda_{j,i}^{-d} \, (d_j^{(j+1)})^d \circ f_{j+1}(X_i^{(j+1)}) = \frac{(1-q_j)^{-d}}{[d]!_{q_j}} \lambda_{j,i}^{-d} \, (d_j^{(j+1)})^d \, (x_i^{(j+1)}) = c_d^{(j+1)}$. So, we get

$$\widetilde{f_{j+1}}(X_i^{(j)}) = x_i^{(j+1)} + \sum_{d=1}^{+\infty} c_d^{(j+1)} (x_j^{(j+1)})^{-d} = x_i^{(j)}$$

So, we always have $\widetilde{f_{j+1}}(X_i^{(j)}) = x_i^{(j)}$ and the restriction f_j of $\widetilde{f_{j+1}}$ to $R^{(j)}$ is the unique homomorphism from $R^{(j)}$ to $A^{(j)}$ which transforms each $X_i^{(j)}$ in $x_i^{(j)}$. f_j is surjective and $ker(f_j) = ker(\widetilde{f_{j+1}}) \cap R^{(j)} = \mathcal{P}^{(j+1)} S_j^{-1} \cap R^{(j)} = \mathcal{P}^{(j)}$. This proves that f_j satisfies the property 3 and it remains to verify that, for any l in $[\![1, \dots, t]\!]$, we have $\overline{h_l} \circ f_j = f_j \circ h_l$. As $h_l(X_i^{(j)}) = \lambda_{l,i} X_i^{(j)}$, it is enough to check that $\overline{h_l}(x_i^{(j)}) = \lambda_{l,i} x_i^{(j)}$ for each i.

If $j \leq i$, we have $\overline{h_l}(x_i^{(j)}) = \overline{h_l}(x_i^{(j+1)}) = \lambda_{l,i} x_i^{(j+1)} = \lambda_{l,i} x_i^{(j)}$.
Now, assume $i < j$.
By the assumption $\mathcal{H}_j$, we can use property 5. at the rank $j+1$, so that

$$\overline{h_l} \circ d_j^{(j+1)} = \lambda_{l,j} d_j^{(j+1)} \circ \overline{h_l}.$$

For each $d \in \mathbb{N}$, we have

$$\overline{h_l}(c_d^{(j+1)}) = \overline{h_l}\Big(\frac{(1-q_j)^{-d}}{[d]!_{q_j}} \lambda_{j,i}^{-d} \, (d_j^{(j+1)})^d \, (x_i^{(j+1)})\Big)$$

$$= \frac{(1-q_j)^{-d}}{[d]!_{q_j}} \lambda_{j,i}^{-d} \, \lambda_{l,j}^{d} (d_j^{(j+1)})^d \circ \overline{h_l}(x_i^{(j+1)}) = \lambda_{l,i} \lambda_{l,j}^d c_d^{(j+1)}.$$

So, $\overline{h_l}(\, c_d^{(j+1)} (x_j^{(j+1)})^{-d}) = \lambda_{l,i} c_d^{(j+1)} (x_j^{(j+1)})^{-d}$, which implies that $\overline{h_l}(x_i^{(j)}) = \lambda_{l,i} x_i^{(j)}$.

□

Recall (see section 3.3) that, since $\mathcal{P}^{(m)}$ is H - invariant, each $\mathcal{P}^{(j)}$ $(2 \leq j \leq m)$ is H - invariant. In particular, $\mathcal{P}^{(2)}$ is H - invariant. Set

$$\Delta = \{ i \in [\![1, \dots, t]\!] \mid Z_i \in \mathcal{P}^{(2)} \} = \{ j_1 < \dots < j_s \} \text{ (unless this set is empty)}$$

and

$$\overline{\Delta} = [\![1, \dots, t]\!] \setminus \Delta = \{ l_1 < \dots < l_e \} \text{ (unless this set is empty).}$$

Recall that $\mathcal{P}^{(2)} = \mathcal{P}_\Delta^{(2)}$ (see section 3.3), $\overline{R} = R^{(2)}$ and set $\overline{A} = A^{(2)}$. For each $i \in [\![1, \dots, t]\!]$, set $z_i = x_i^{(2)}$ and observe that $f_2 : \overline{R} \to \overline{A}$ transforms each $Z_i = X_i^{(2)}$ in z_i. So, we can also describe Δ and $\overline{\Delta}$ as follows:

$$\Delta = \{ i \in [\![1, \dots, t]\!] \mid z_i = 0 \}$$

$$\overline{\Delta} = \{ i \in [\![1, \dots, t]\!] \mid z_i \neq 0 \}$$

Observation 4.2. 1
Assume that $\overline{\Delta}$ is empty. Then each $x_i^{(m)}$ $(i \in [\![1, \dots, t]\!])$ is zero.

Proof
If $\overline{\Delta}$ is empty, we have $x_j^{(j+1)} = z_j = 0$ for each $j \in [\![1, \dots, m-1]\!]$. By (proposition 4.2. 1, 6.), this implies that, if $i \in [\![1, \dots, t]\!]$ and $j \in [\![1, \dots, m-1]\!]$, we have $x_i^{(j+1)} = x_i^{(j)}$. So, each $x_i^{(m)} = x_i^{(2)} = z_i = 0$.

□

Until the end of section 4.2, we assume that $\overline{\Delta}$ is nonempty. So, we have:

Proposition 4.2. 2

1. $\overline{A} = k < z_{l_1}, \dots, z_{l_e} >$.
2. *If* $1 \leq i < d \leq e$, *we have* $z_{l_d} z_{l_i} = \lambda_{l_d, l_i} z_{l_i} z_{l_d}$.
3. $z_{l_1}, \dots, z_{l_e}$ *are all nonzero and the Laurent (ordered) monomials* $z^{\underline{a}} = z_{l_1}^{a_1} \dots z_{l_e}^{a_e}$ $(\underline{a} = (a_1, \dots, a_e) \in \mathbb{Z}^e)$ *are* k *- linearly independent.*

Proof.

1. $\overline{A} = k < z_1, \dots, z_t >$ (see proposition 4.2. 1) and, for $i \notin \{l_1, \dots, l_e\}$, we have $i \in \Delta \Rightarrow Z_i \in \mathcal{P}^{(2)} = ker(f_2) \Rightarrow z_i = 0$.
2. We know (section 3.2) that $Z_{l_d} Z_{l_i} = \lambda_{l_d, l_i} Z_{l_i} Z_{l_d}$. If we transform by f_2, we obtain the required equality.
3. Denote by $\widehat{R} = k < Z_{l_1}, \dots, Z_{l_e} >$ the subalgebra of $\overline{R}$ generated by $Z_{l_1}, \dots, Z_{l_e}$. By the property 1., f_2 induces (by restriction) a surjective homomorphism $\widehat{f_2} : \widehat{R} \to \overline{A}$ and $ker(\widehat{f_2}) = \widehat{R} \cap \mathcal{P}^{(2)}$. As each $P \in \mathcal{P}^{(2)} = \mathcal{P}_{\Delta}^{(2)} = (\{Z_{j_1}, \dots, Z_{j_s}\})$ is a linear combination of monomials in which at least one of the variables Z_{j_i} with $1 \leq i \leq s$ appears, this intersection is reduced to zero and then, $\widehat{f_2}$ is an isomorphism which transforms each Z_{l_i} in z_{l_i}. As the ordered monomials in $Z_{l_1}, \dots, Z_{l_e}$ are linearly independent, we have the same property for the ordered monomials in $z_{l_1}, \dots, z_{l_e}$. This easily implies that $z_{l_1}, \dots, z_{l_e}$ are nonzero and that the Laurent ordered monomials in $z_{l_1}, \dots, z_{l_e}$ are also linearly independent.

□

4.3 Each $x_i^{(m)}$ is a Laurent polynomial in $z_{l_1}, \dots, z_{l_e}$.

The conventions are the same as in section 4.2. and we still assume that $\overline{\Delta}$ is nonempty.

Let us consider some i in $[\![1, \dots, t]\!]$.

If $u \in D_m$ and if $\gamma \in \mathbb{Z}\Pi$, we say (as in section 3.2) that u *is homogeneous of degree* γ if $\overline{h_\rho}(u) = q^{-(\rho,\gamma)} u$ for all ρ in $\mathbb{Z}\Pi$. Since q is not a root of unity and V is spanned by Π, the degree of a nonzero homogeneous element is uniquely defined. Moreover, we immediately have the following properties:

- If $u_1, \dots, u_r$ are homogeneous of same degree γ, then any linear combination of $u_1, \dots, u_r$ (with coefficients in k) is homogeneous of degree γ.
- If u_1 is homogeneous of degree γ_1 and u_2 is homogeneous of degree γ_2, then $u_1 u_2$ is homogeneous of degree $\gamma_1 + \gamma_2$.
- Likewise, if u is nonzero and homogeneous of degree γ, then u^{-1} is homogeneous of degree $-\gamma$.
- So, if $u_1, \dots, u_r$ are nonzero and homogeneous of degrees $\gamma_1, \dots, \gamma_r$ respectively, if $\underline{a} = (a_1, \dots, a_r) \in \mathbb{Z}^r$, then $u^{\underline{a}} := u_1^{a_1} \dots u_r^{a_r}$ is homogeneous of degree $a_1\gamma_1 + \dots + a_r\gamma_r$.

Lemma 4.3. 1
Consider some j *in* $[\![2, \ldots , m]\!]$.

1. *If* $j \leq i+1$, *then* $x_i^{(j)} = z_i$. *In particular,* $x_i^{(i+1)} = z_i$.
2. *If* $1 \leq i < l \leq t$, *then* $x_l^{(j)} x_i^{(j)} - \lambda_{l,i} x_i^{(j)} x_l^{(j)} = d_l^{(j)}(x_i^{(j)}) = f_j(P_{l,i}^{(j)}) \in k < x_{i+1}^{(j)}, \ldots , x_{l-1}^{(j)} >$. *Moreover, if* $l \geq j$ *or* $l = i+1$, *then* $d_l^{(j)}(x_i^{(j)}) = 0$.
3. *If* $U \in R^{(j)}$ *is homogeneous of degree* γ, *then* $f_j(U)$ *is homogeneous of same degree* γ. *In particular,* $x_i^{(j)}$ *is homogeneous of degree* β_i.
4. *If* $j \leq l \leq t$ *and* $u \in k < x_1^{(j)}, \ldots , x_{l-1}^{(j)} >$, *then* $z_l u = \overline{h_l}(u) z_l$.

Proof

1. By proposition 4.2. 1, we have $x_i^{(i+1)} = x_i^{(i)} = \ldots = x_i^{(2)} = z_i$.
2. By proposition 4.2. 1, we have $x_l^{(j)} x_i^{(j)} - \lambda_{l,i} x_i^{(j)} x_l^{(j)} = x_l^{(j)} x_i^{(j)} - \overline{h_l}(x_i^{(j)}) x_l^{(j)} = d_l^{(j)}(x_i^{(j)}) = d_l^{(j)} \circ f_j(X_i^{(j)}) = f_j \circ \delta_l^{(j)}(X_i^{(j)})$ and we know (see section 3.2) that $\delta_l^{(j)}(X_i^{(j)}) = P_{l,i}^{(j)} \in k < X_{i+1}^{(j)}, \ldots , X_{l-1}^{(j)} >$. Moreover, if $l \geq j$ or $l = i+1$, we know (see section 3.2) that $P_{l,i}^{(j)} = 0$. Transforming by f_j, we get the required result.
3. Consider ρ in $\mathbb{Z}\Pi$. By proposition 4.2. 1, we have $\overline{h_\rho}(f_j(U)) = f_j(h_\rho(U)) = q^{-(\rho,\gamma)} f_j(U)$.
4. By 1., we have $z_l = x_l^{(j)}$. So, $z_l u - \overline{h_l}(u) z_l = x_l^{(j)} u - \overline{h_l}(u) x_l^{(j)} = d_l^{(j)}(u)$.
   By 2., if $i < l$, we have $d_l^{(j)}(x_i^{(j)}) = 0$ (since $j \leq l$). This implies that $d_l^{(j)}$ is zero on $k < x_1^{(j)}, \ldots , x_{l-1}^{(j)} >$ and, in particular, $d_l^{(j)}(u) = 0$.

□

Lemma 4.3. 2
Consider some j *in* $[\![2, \ldots , m]\!]$.

1. *If* $j \leq l_1$, *then* $x_i^{(j)} = z_i$.
2. *Assume that* $l_1 < j$ *and denote by* d *the greatest integer such that* $l_d < j$ $(1 \leq d \leq p)$. *Then* $x_i^{(j)} = x_i^{(l_d+1)}$.

Proof

2. We prove this by induction on j. If $j = l_d + 1$, there is nothing to prove. Assume $j > l_d + 1$ and set $j' = j - 1 > l_d$. By lemma 4.3. 1, we have $x_{j'}^{(j'+1)} = z_{j'} = 0$ (since $j' \in \Delta$). So, by proposition 4.2. 1, $x_i^{(j)} = x_i^{(j'+1)} = x_i^{(j')} = x_i^{(l_d+1)}$ (by the induction assumption).

1. The proof is the same (observing that there is nothing to prove if $j = 2$).

□

Now, assume that $l_1 < m$ and denote by p the greatest integer such that $l_p < m$.

Proposition 4.3. 1
Consider some j *in* $[\![2, \ldots , m]\!]$ *and assume that* $i < j$.

1. *If* $i < j \leq l_1$, *then* $x_i^{(j)} = z_i = 0$.

   Assume that $l_1 < j$ *and denote by* d *the greatest integer such that* $l_d < j$ $(1 \leq d \leq p)$.

2. *(a) If* $l_d \leq i < j$, *then* $x_i^{(j)} = z_i$.

   (b) If $i < l_d < j$, *then*

$$x_i^{(j)} = x_i^{(l_d)} + Q_1 z_{l_d}^{-1} + \ ... \ + Q_K z_{l_d}^{-K}$$

   with $K \geq 1$ *and:*

   - *If* $d = 1$, *then each* $Q_l \in k$.
   - *If* $d > 1$ *and* $i \geq l_{d-1}$, *then each* $Q_l \in k$.
   - *If* $d > 1$ *and* $i < l_{d-1}$, *then each* $Q_l \in k < x_{i+1}^{(l_d)}, \ ... \ , x_{l_{d-1}}^{(l_d)} >$.

Proof

1. By (lemma 4.3. 2, 1.), we have $x_i^{(j)} = z_i$ and, since $i < l_1$, we have $i \in \Delta$, so that $z_i = 0$.

2. (a) By (lemma 4.3. 2, 2.), we have $x_i^{(j)} = x_i^{(l_d+1)}$ and, since $l_d + 1 \leq i$, this is also equal to z_i.

   (b) We prove this result by decreasing induction on i.
   As in 2.a., we have $x_i^{(j)} = x_i^{(l_d+1)}$ and, since $x_{l_d}^{(l_d+1)} = z_{l_d} \neq 0$, it results from (proposition 4.2. 1, 7.(b)) that

$$x_i^{(j)} = x_i^{(l_d)} + P_1 z_{l_d}^{-1} + \ ... \ + P_L z_{l_d}^{-L}$$

   with $L \geq 1$ and each $P_l = \mu_l \ (d_{l_d}^{(l_d+1)})^l \ (x_i^{(l_d+1)}) \ (\mu_l \in k^*)$. So, by lemma 4.3. 1, each $P_l \in k < x_{i+1}^{(l_d+1)}, \ ... \ , x_{l_d-1}^{(l_d+1)} > = k < x_{i+1}^{(j)}, \ ... \ , x_{l_d-1}^{(j)} >$ (by lemma 4.3. 2).

   Assume that $i = l_d - 1$. Then, by lemma 4.3. 1, we have $d_{l_d}^{(l_d+1)} \ (x_i^{(l_d+1)}) = 0$. This implies that $P_l = 0$ for each l. So, $x_i^{(j)} = x_i^{(l_d)}$ and the proof is over.

   Now, assume that $i < l_d - 1$ and, for $i + 1 \leq h \leq l_d - 1$,

$$x_h^{(j)} = x_h^{(l_d)} + Q_{1,h} z_{l_d}^{-1} + \ ... \ + Q_{M,h} z_{l_d}^{-M}$$

   with $M \geq 1$, and:

   - each $Q_{l,h} \in k$ if $d = 1$,
   - $Q_{l,h} \in k$ if $d > 1$ and $h \geq l_{d-1}$,
   - $Q_{l,h} \in k < x_{h+1}^{(l_d)}, \ ... \ x_{l_{d-1}}^{(l_d)} >$ if $d > 1$ and $h < l_{d-1}$.

   This implies the following results:

   - If $d = 1$, then each $x_h^{(j)} \in k < x_h^{(l_d)}, z_{l_d}^{-1} >$ $(i + 1 \leq h \leq l_d - 1)$.
   - If $d > 1$ and $l_{d-1} \leq h \leq l_d - 1$, then $x_h^{(j)} \in k < x_h^{(l_d)}, z_{l_d}^{-1} >$.
   - If $d > 1$ and $i + 1 \leq h < l_{d-1}$, then $x_h^{(j)} \in k < x_h^{(l_d)}, x_{h+1}^{(l_d)}, \ ... \ , x_{l_{d-1}}^{(l_d)}, z_{l_d}^{-1} >$.

   So, it turns out that each $P_l \in k < x_{i+1}^{(l_d)}, \ ... \ , x_{l_d-1}^{(l_d)}, z_{l_d}^{-1} >$.

   - If $d = 1$ and $h < l_d$, we have (by the first point of the proposition) $x_h^{(l_d)} = z_h = 0$. So, in this case, each $P_l \in k < z_{l_d}^{-1} >$ and

$$x_i^{(j)} = x_i^{(l_d)} + Q_1 z_{l_d}^{-1} + \ ... \ + Q_K z_{l_d}^{-K}$$

     with $K \geq 1$ and each Q_l in k.
   - Now, assume that $d > 1$.
     If $l_{d-1} < h \leq l_d - 1$, we have (by 2.a.) $x_h^{(l_d)} = z_h = 0$ (since $h \in \Delta$). So,
     - If $i \geq l_{d-1}$, each $P_l \in k < z_{l_d}^{-1} >$ and

$$x_i^{(j)} = x_i^{(l_d)} + Q_1 z_{l_d}^{-1} + \ ... \ + Q_K z_{l_d}^{-K}$$

       with $K \geq 1$ and each Q_l in k.

– If $i < l_{d-1}$, each $P_l \in k < x_{i+1}^{(l_d)}, \ldots, x_{l_{d-1}}^{(l_d)}, z_{l_d}^{-1} >$. As, by lemma 4.3. 1, $z_{l_d} = x_{l_d}^{(l_d)}$ q-commutes with $x_{i+1}^{(l_d)}, \ldots, x_{l_{d-1}}^{(l_d)}$, each P_l can be written as follows:

$$P_l = S_{0,l} + S_{1,l} z_{l_d}^{-1} + \ldots + S_{E,l} z_{l_d}^{-E}$$

with $E \geq 1$ and each $S_{j,l} \in k < x_{i+1}^{(l_d)}, \ldots, x_{l_{d-1}}^{(l_d)} >$. So, we conclude that, in this case,

$$x_i^{(j)} = x_i^{(l_d)} + Q_1 z_{l_d}^{-1} + \ldots + Q_K z_{l_d}^{-K}$$

with $K \geq 1$ and each Q_l in $k < x_{i+1}^{(l_d)}, \ldots, x_{l_{d-1}}^{(l_d)} >$.

□

Proposition 4.3. 2

Consider some j *in* $[\![2, \ldots, m]\!]$, *assume that* $i+1 < j$ *and consider some* $u \in k < x_{i+1}^{(j)}, \ldots, x_{j-1}^{(j)} >$.

1. *If* $j \leq l_1$, *then* $u \in k$.
   Assume that $l_1 < j$ *and denote by* d *the greatest integer such that* $l_d < j$ $(1 \leq d \leq p)$.
2. (a) *If* $l_d \leq i$, *then* $u \in k$.
   (b) *If* $i < l_d$, *then*
   $$u = u_1 z_{l_d}^{a_1} + \ldots + u_K z_{l_d}^{a_K}$$
   with $K \geq 1$, $(a_1, \ldots, a_K) \in \mathbb{Z}^K$ *and:*
   - *If* $d = 1$, *then each* $u_l \in k$.
   - *If* $d > 1$ *and* $i \geq l_{d-1}$, *then each* $u_l \in k$.
   - *If* $d > 1$ *and* $i < l_{d-1}$, *then each* $u_l \in k < x_{i+1}^{(l_d)}, \ldots, x_{l_{d-1}}^{(l_d)} >$.

Proof

1. By (proposition 4.3. 1, 1.) we have $x_h^{(j)} = 0$ for any $h < j$. So, $u \in k$.
2. (a) Consider some integer h such that $l_d < h < j$. By (proposition 4.3. 1, 2.a.) we have $x_h^{(j)} = z_h = 0$ (since $h \in \Delta$). So, $u \in k$.
   (b) Consider some integer h with $i < h < j$.
   If $l_d < h$, then, as in 2.a., we have $x_h^{(j)} = 0$. If $h = l_d$, then, again by (proposition 4.3. 1, 2.a.), we have $x_h^{(j)} = z_h = z_{l_d}$.
   - Assume that $d = 1$. If $h < l_d$, then, by (proposition 4.3. 1, 2.b.), we have
     $$x_h^{(j)} = x_h^{(l_d)} + Q_1 z_{l_d}^{-1} + \ldots + Q_M z_{l_d}^{-M}$$
     with $M \geq 1$ and each Q_l in k. By (proposition 4.3. 1, 1.), we also have $x_h^{(l_d)} = 0$. So, $x_h^{(j)} \in k < z_{l_d}^{-1} >$ and it turns out that $u \in k < z_{l_d}^{\pm 1} >$. This implies that
     $$u = u_1 z_{l_d}^{a_1} + \ldots + u_K z_{l_d}^{a_K}$$
     with $N \geq 1$, $(a_1, \ldots, a_N) \in \mathbb{Z}^N$ and each $u_l \in k$.
   - Assume that $d > 1$ and $i \geq l_{d-1}$. If $i < h < l_d$, we have $l_{d-1} < h < l_d$ and, by (proposition 4.3. 1, 2.b.),
     $$x_h^{(j)} = x_h^{(l_d)} + Q_1 z_{l_d}^{-1} + \ldots + Q_M z_{l_d}^{-M}$$
     with $M \geq 1$ and each $Q_l \in k$. By (proposition 4.3. 1, 1.) we also have $x_h^{(l_d)} = z_h$ and, since $h \in \Delta$, $x_h^{(l_d)} = 0$. As over, we conclude that
     $$u = u_1 z_{l_d}^{a_1} + \ldots + u_K z_{l_d}^{a_K}$$
     with $K \geq 1$, $(a_1, \ldots, a_K) \in \mathbb{Z}^K$ and each $u_l \in k$.

- Assume that $d > 1$ and $i < l_{d-1}$. If $i < h < l_d$, we have, always by (proposition 4.3. 1, 2.b.),

$$x_h^{(j)} = x_h^{(l_d)} + Q_1 z_{l_d}^{-1} + \dots + Q_M z_{l_d}^{-M}$$

with $M \geq 1$ and:
  - If $l_{d-1} < h$, then each $Q_l \in k$ and, as in the previous case, $x_h^{(l_d)} = 0$. So, $x_h^{(j)} \in k < z_{l_d}^{-1} >$.
  - If $h = l_{d-1}$, then each $Q_l \in k$ and, consequently, $x_h^{(j)} \in k < x_h^{(l_d)}, z_{l_d}^{-1} > = k < x_{l_{d-1}}^{(l_d)}, z_{l_d}^{-1} >$.
  - If $h < l_{d-1}$, then each $Q_l \in < x_{h+1}^{(l_d)}, \dots , x_{l_{d-1}}^{(l_d)} >$ and, consequently, $x_h^{(j)} \in k < x_h^{(l_d)}, x_{h+1}^{(l_d)}, \dots , x_{l_{d-1}}^{(l_d)}, z_{l_d}^{-1} >$.
  So, it turns out that $u \in k < x_{i+1}^{(l_d)}, \dots , x_{l_{d-1}}^{(l_d)}, z_{l_d}^{\pm 1} >$. As, by lemma 4.3. 1, $z_{l_d} = x_{l_d}^{(l_d)}$ q-commutes with $x_{i+1}^{(l_d)}, \dots , x_{l_{d-1}}^{(l_d)}$, we conclude that

$$u = u_1 z_{l_d}^{a_1} + \dots + u_K z_{l_d}^{a_K}$$

  with $K \geq 1$, $(a_1, \dots , a_K) \in \mathbb{Z}^K$ and each u_l in $k < x_{i+1}^{(l_d)}, \dots , x_{l_{d-1}}^{(l_d)} >$.

□

Proposition 4.3. 3
Consider some j *in* $[\![2, \dots , m]\!]$ *with* $i < j$.

1. *If* $\{l_1, \dots , l_p\} \cap [\![i+1, \dots , j-1]\!] = \emptyset$, *then* $x_i^{(j)} = z_i$.
2. *Assume that* $\{l_1, \dots , l_p\} \cap [\![i+1, \dots , j-1]\!] \neq \emptyset$ *and set* $\{l_1, \dots , l_p\} \cap [\![i+1, \dots , j-1]\!] = \{l_c < \dots < l_d\}$. *Then, we have*

$$x_i^{(j)} = z_i + \sum_{\underline{a} = (a_c, \dots , a_d) \in F} \eta(\underline{a}) z_{l_c}^{a_c} \dots z_{l_d}^{a_d},$$

   where
   - F *is a finite (possibly empty) subset of* $\mathbb{Z}^{d-c+1}$.
   - *If* $\preceq$ *denotes the inverse lexicographic order in* $\mathbb{Z}^{d-c+1}$, *then, for any* $\underline{a}$ *in* F, *we have* $\underline{a} \prec 0$ *(ie.* $\underline{a} = (a_c, \dots , a_r, 0, \dots , 0)$ *with* $a_r < 0$.)
   - *For each* $\underline{a}$ *in* F, *we have* $\eta(\underline{a}) \in k^*$.

Proof
We prove the proposition by induction on $j - i$. In order to do this, we first prove the assertion 1.

So, assume that $\{l_1, \dots , l_p\} \cap [\![i+1, \dots , j-1]\!] = \emptyset$, and observe that there are two possibilities:

1. (a) $j \leq l_1$. By proposition 4.3. 1, we know that $x_i^{(j)} = z_i$.
   (b) $l_1 < j$ and, if d is the greatest integer such that $l_d < j$, $l_d \leq i < j$. Again, by proposition 4.3. 1, we know that $x_i^{(j)} = z_i$.

   Assume $j - i = 1$. In this case, we have $\{l_1, \dots , l_p\} \cap [\![i+1, \dots , j-1]\!] = \emptyset$. So, we are in the case of assertion 1., and the proof is over.

Now assume $j - i > 1$. As the assertion 1. is already proved, we may assume that $\{l_1, \dots , l_p\} \cap [\![i+1, \dots, j-1]\!] = \{l_c < \dots < l_d\}$ is non empty. So, we have $l_1 < j$, d is the greatest integer such that $l_d < j$ and $i < l_d < j$. Now, by proposition 4.3. 1, we can write

$$x_i^{(j)} = x_i^{(l_d)} + Q_1 z_{l_d}^{-1} + \dots + Q_K z_{l_d}^{-K}$$

with $K \geq 1$ and three possible cases:

1. $d = 1$. Then each $Q_l \in k$ and, since $x_i^{(l_d)} = z_i$ (see proposition 4.3. 1, 1.), the proof is over.

2. $d > 1$ and $i \geq l_{d-1}$. Then each $Q_l \in k$ and, since $x_i^{(l_d)} = z_i$ (see proposition 4.3. 1, 2.a.), the proof is over.
3. $d > 1$ and $i < l_{d-1}$ (so that $c \leq d-1$). In this case, each $Q_l \in k < x_{i+1}^{(l_d)}, \ \dots \ , \ x_{l_{d-1}}^{(l_d)} >$. Since $l_d < j$, it results from the induction assumption that the proposition is true for each $x_h^{(l_d)}$ with $i \leq h < l_d$. This implies that
   - If $i+1 \leq h \leq l_{d-1}$, we have $x_h^{(l_d)} \in k < z_{l_c}^{\pm 1}, \ \dots \ , \ z_{l_{d-1}}^{\pm 1} >$ (since $\{l_1, \ \dots \ , \ l_p\} \ \cap \ \llbracket h+1, \dots, l_d - 1 \rrbracket \subseteq \{l_c \ < \dots < \ l_{d-1}\}$ and z_h is either 0 or equal to z_{l_r} with $c \leq r \leq d-1$). So, each $Q_l \in k < z_{l_c}^{\pm 1}, \ \dots \ , \ z_{l_{d-1}}^{\pm 1} >$.
   - $$x_i^{(l_d)} \ = \ z_i \ + \sum_{\underline{a} \ = \ (a_c, \ \dots \ , \ a_{d-1}) \ \in \ F_1} \eta(\underline{a}) z_{l_c}^{a_c} \ \dots \ z_{l_{d-1}}^{a_{d-1}},$$

     where
     - ◇ F_1 is a finite (possibly empty) subset of $\mathbb{Z}^{d-c}$.
     - ◇ For each $\underline{a}$ in F_1, we have $\underline{a} \ \prec \ 0$ and $\eta(\underline{a}) \ \in \ k^*$.

   As each Q_l is a summand of (ordered) Laurent monomials in $z_{l_c}, \ \dots \ , \ z_{l_{d-1}}$, it turns out that

   $$x_i^{(j)} \ = \ z_i \ + \sum_{\underline{a} \ = \ (a_c, \ \dots \ , \ a_d) \ \in \ F} \eta(\underline{a}) z_{l_c}^{a_c} \ \dots \ z_{l_d}^{a_d},$$

   where
   - ◇ F is a finite (possibly empty) subset of $\mathbb{Z}^{d-c+1}$.
   - ◇ For each $\underline{a}$ in F, we have $\underline{a} \ \prec \ 0$ and $\eta(\underline{a}) \ \in \ k^*$.

□

This proposition implies immediately

Corollary 4.3. 1 *Consider some integer* d *with* $1 < d \leq p$. *If* $1 \leq i < l_d$, *then* $x_i^{(l_d)} \ \in \ k < z_{l_1}^{\pm 1}, \ \dots \ , \ z_{l_{d-1}}^{\pm 1} >$.

Proof
By proposition 4.3. 3, $x_i^{(l_d)} \ = \ z_i + Q$ with Q a (possibly 0) Laurent polynomial in $z_{l_1}, \ \dots \ , \ z_{l_{d-1}}$. As z_i is either 0 or z_{l_r} with $1 \leq r \leq d-1$, the proof is over.

□

Corollary 4.3. 2 *Consider some integer* d *with* $1 < d \leq p$. *In* D_m *considered as a left module over* $k < x_1^{(l_d)}, \ \dots \ , \ x_{l_d - 1}^{(l_d)} >$, *the ordered Laurent monomials* $z_{l_d}^{a_d} \ \dots \ z_{l_p}^{a_p}$ ($a_d, \ \dots \ , \ a_p$ *in* $\mathbb{Z}$) *are linearly independent.*

Proof
By corollary 4.3. 1, we have $k < x_1^{(l_d)}, \ \dots \ , \ x_{l_d - 1}^{(l_d)} > \ \subseteq \ k < z_{l_1}^{\pm 1}, \ \dots \ , \ z_{l_{d-1}}^{\pm 1} >$. So, the corollary results immediately from the k - linear independence of the ordered Laurent monomials in $z_{l_1}, \ \dots \ , \ z_{l_e}$ (see proposition 4.2. 2).

□

4.4 A new sufficient condition for $\mathcal{P}^{(m)}$ to be in $Im(\phi_m)$.

The notations are the same as in the preceding sections, but we do not assume that $\overline{\Delta}$ is nonempty a priori.
Assume that $\mathcal{P}^{(m)}$ is not in $Im(\phi_m)$ so that, by proposition 4.1. 1, $X_m^{(m)} \ \in \ \mathcal{P}^{(m)}$ and there is some j in $\llbracket 1, \ \dots \ , m-1 \rrbracket$ such that

$$U \ = \ \Theta^{(m)}(\delta_m^{(m+1)}(X_j^{(m+1)})) \ \notin \ \mathcal{P}^{(m)}.$$

Let us choose such a j maximal. So, we have

$$\Theta^{(m)}(\delta_m^{(m+1)}(X_i^{(m+1)})) \ \in \ \mathcal{P}^{(m)}$$

for any i in $\llbracket j+1, \ \dots \ , m-1 \rrbracket$ and we observe that, since $U \ \notin \ \mathcal{P}^{(m)}$, we have $u := f_m(U) \neq 0$ in $A^{(m)}$.

Recall that

$$\Delta \ = \ \{i \in \ \llbracket 1, \ \dots \ , \ t \rrbracket \mid Z_i \in \ \mathcal{P}^{(2)}\} \ = \ \{j_1 \ < \ \dots \ < \ j_s\} \text{ (unless this set is empty)},$$
$$\overline{\Delta} \ = \ \llbracket 1, \ \dots \ , \ t \rrbracket \ \setminus \ \Delta \ = \ \{l_1 \ < \ \dots \ < \ l_e\} \text{ (unless this set is empty)}.$$

Lemma 4.4. 1
$\overline{\Delta}$ is nonempty, $l_1 < m$, $m \notin \overline{\Delta}$ and, if p denotes the greatest integer such that $l_p < m$, we have $j < l_p < m$.

Proof
First, recall that:

- If $\overline{\Delta}$ is empty, then (observation 4.2. 1) each $x_l^{(m)}$ ($l \in \llbracket 1, \dots , t \rrbracket$) is zero.
- $\delta_m^{(m+1)}(X_j^{(m+1)}) = P_{m,j}^{(m+1)}$. This is 0 if $m = j+1$ or a (finite) summand
$$\sum_{\underline{a} \, = \, (a_{j+1}, \dots, a_{m-1})} c_{\underline{a}} (X_{j+1}^{(m+1)})^{a_{j+1}} \dots (X_{m-1}^{(m+1)})^{a_{m-1}}$$
with each $\underline{a}$ in $\mathbb{N}^{m-1-j}$ if $m > j+1$ and, moreover, $\underline{a}$ is nonzero when $c_{\underline{a}} \neq 0$.
- $\Theta^{(m)}$: $k < X_1^{(m+1)}, \dots , X_{m-1}^{(m+1)} > \ \to \ k < X_1^{(m)}, \dots , X_{m-1}^{(m)} >$ is an algebra homomorphism which transforms each $X_l^{(m+1)}$ in $X_l^{(m)}$.

Assume that $m = j+1$. In this case, we have

$$\delta_m^{(m+1)}(X_j^{(m+1)}) = 0 \ \Rightarrow \ U = \Theta^{(m)}(\delta_m^{(m+1)}(X_j^{(m+1)})) = 0 \ \Rightarrow \ u = f_m(U) = 0$$

and we obtain a contradiction.

So, we have $m > j+1$, and

$$\delta_m^{(m+1)}(X_j^{(m+1)}) = \sum_{\underline{a} \, = \, (a_{j+1}, \dots, a_{m-1})} c_{\underline{a}} (X_{j+1}^{(m+1)})^{a_{j+1}} \dots (X_{m-1}^{(m+1)})^{a_{m-1}} \ \Rightarrow$$

$$U = \Theta^{(m)}(\delta_m^{(m+1)}(X_j^{(m+1)})) = \sum_{\underline{a} \, = \, (a_{j+1}, \dots, a_{m-1})} c_{\underline{a}} (X_{j+1}^{(m)})^{a_{j+1}} \dots (X_{m-1}^{(m)})^{a_{m-1}} \ \Rightarrow$$

$$u = f_m(U) = \sum_{\underline{a} \, = \, (a_{j+1}, \dots, a_{m-1})} c_{\underline{a}} (x_{j+1}^{(m)})^{a_{j+1}} \dots (x_{m-1}^{(m)})^{a_{m-1}}.$$

Assume that $\overline{\Delta}$ is empty. Then, each $x_l^{(m)}$ is zero and, since $\underline{a}$ is nonzero when $c_{\underline{a}} \neq 0$, we get that $u = 0$ and we still have a contradiction. So, $\overline{\Delta}$ is nonempty.

If $l_1 \geq m$, then, for each $l \in \llbracket 1, \dots , m-1 \rrbracket$, we have (by lemma 4.3. 2 and, since $l < l_1$) $x_l^{(m)} = z_l = 0$. As over, this implies that $u = 0$ and we still get a contradiction . So, $l_1 < m$.

Since $X_m^{(m)} \in \mathcal{P}^{(m)}$, we have $x_m^{(m)} = 0$. As $x_m^{(m)} = z_m$ (lemma 4.3. 1), we have $Z_m \in \mathcal{P}^{(2)}$ and so, $m \in \Delta$. Moreover, if $l_p \leq j$, then, for each $l \in \llbracket j+1, \dots , m-1 \rrbracket$, we have (by propostion 4.3. 1, 2.a. and, since $l_p < l < m$) $x_l^{(m)} = z_l = 0$. As over, this implies that $u = 0$ and we still get a contradiction . So, $j < l_p$.

□

Now, set $V = \delta_m^{(m+1)}(X_j^{(m+1)}) = X_m^{(m+1)} X_j^{(m+1)} - q^{-(\beta_m, \beta_j)} X_j^{(m+1)} X_m^{(m+1)}$ (see section 3.2), so that $U = \Theta^{(m)}(V)$.

Lemma 4.4. 2
Consider some integer l with $j < l < m$. Then we have, in $R^{(m+1)}$,

$$X_l^{(m+1)} V = q^{(\beta_l, \beta_m - \beta_j)} V X_l^{(m+1)}$$

$$+ \ q^{(\beta_l, \beta_m)} \left[\ q^{-(\beta_j, \beta_l + \beta_m)} X_j^{(m+1)} \delta_m^{(m+1)}(X_l^{(m+1)}) - \delta_m^{(m+1)}(X_l^{(m+1)}) X_j^{(m+1)} \ \right]$$

$$+ \ q^{(\beta_l, \beta_m)} \delta_m^{(m+1)} \circ \delta_l^{(m+1)}(X_j^{(m+1)}).$$

Proof
We observe that, as each $X_i^{(m+1)}$ is homogeneous of degree β_i (see section 3.2), we have

$$\delta_l^{(m+1)}(V) = \delta_l^{(m+1)}(X_m^{(m+1)})X_j^{(m+1)} + q^{-(\beta_l,\beta_m)}X_m^{(m+1)}\delta_l^{(m+1)}(X_j^{(m+1)})$$
$$-q^{-(\beta_m,\beta_j)}\delta_l^{(m+1)}(X_j^{(m+1)})X_m^{(m+1)} - q^{-[(\beta_m,\beta_j)+(\beta_l,\beta_j)]}X_j^{(m+1)}\delta_l^{(m+1)}(X_m^{(m+1)}).$$

So, we can write

$$\delta_l^{(m+1)}(V) = A + B - C - D$$

with

$$A = \delta_l^{(m+1)}(X_m^{(m+1)})X_j^{(m+1)},$$
$$B = q^{-(\beta_l,\beta_m)}X_m^{(m+1)}\delta_l^{(m+1)}(X_j^{(m+1)}),$$
$$C = q^{-(\beta_m,\beta_j)}\delta_l^{(m+1)}(X_j^{(m+1)})X_m^{(m+1)},$$
$$D = q^{-[(\beta_m,\beta_j)+(\beta_l,\beta_j)]}X_j^{(m+1)}\delta_l^{(m+1)}(X_m^{(m+1)}).$$

Now, we compute separately the different pieces of the right member:

- $\delta_l^{(m+1)}(X_m^{(m+1)}) = X_l^{(m+1)}X_m^{(m+1)} - q^{-(\beta_l,\beta_m)}X_m^{(m+1)}X_l^{(m+1)}$. As we also have $\delta_m^{(m+1)}(X_l^{(m+1)}) = X_m^{(m+1)}X_l^{(m+1)} - q^{-(\beta_l,\beta_m)}X_l^{(m+1)}X_m^{(m+1)}$, we can write $X_l^{(m+1)}X_m^{(m+1)} = q^{(\beta_l,\beta_m)}\ [\ X_m^{(m+1)}X_l^{(m+1)} - \delta_m^{(m+1)}(X_l^{(m+1)})\ ]$.
  So, we get:

  $$\delta_l^{(m+1)}(X_m^{(m+1)}) = (q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)})X_m^{(m+1)}X_l^{(m+1)} - q^{(\beta_l,\beta_m)}\delta_m^{(m+1)}(X_l^{(m+1)}).$$

- From this, we deduce that

  $$\delta_l^{(m+1)}(X_m^{(m+1)})X_j^{(m+1)} = (q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)})X_m^{(m+1)}X_l^{(m+1)}X_j^{(m+1)} - q^{(\beta_l,\beta_m)}\delta_m^{(m+1)}(X_l^{(m+1)})X_j^{(m+1)}.$$

  But we also have

  $\delta_l^{(m+1)}(X_j^{(m+1)}) = X_l^{(m+1)}X_j^{(m+1)} - q^{-(\beta_l,\beta_j)}X_j^{(m+1)}X_l^{(m+1)}$, so that $\delta_l^{(m+1)}(X_m^{(m+1)})X_j^{(m+1)} =$ $(q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)})X_m^{(m+1)}[q^{-(\beta_l,\beta_j)}X_j^{(m+1)}X_l^{(m+1)} + \delta_l^{(m+1)}(X_j^{(m+1)})] - q^{(\beta_l,\beta_m)}\delta_m^{(m+1)}(X_l^{(m+1)})X_j^{(m+1)}$.

  So, we get:

  $$A = (q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)})q^{-(\beta_l,\beta_j)}X_m^{(m+1)}X_j^{(m+1)}X_l^{(m+1)}$$
  $$+ (q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)})X_m^{(m+1)}\delta_l^{(m+1)}(X_j^{(m+1)})$$
  $$- q^{(\beta_l,\beta_m)}\delta_m^{(m+1)}(X_l^{(m+1)})X_j^{(m+1)}.$$

- Now, we have $A + B - C =$
  $(q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)})q^{-(\beta_l,\beta_j)}X_m^{(m+1)}X_j^{(m+1)}X_l^{(m+1)}$
  $+ q^{(\beta_l,\beta_m)}X_m^{(m+1)}\delta_l^{(m+1)}(X_j^{(m+1)}) - q^{-(\beta_m,\beta_j)}\delta_l^{(m+1)}(X_j^{(m+1)})X_m^{(m+1)}$
  $- q^{(\beta_l,\beta_m)}\delta_m^{(m+1)}(X_l^{(m+1)})X_j^{(m+1)} =$
  $(q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)})q^{-(\beta_l,\beta_j)}X_m^{(m+1)}X_j^{(m+1)}X_l^{(m+1)}$
  $+ q^{(\beta_l,\beta_m)}[X_m^{(m+1)}\delta_l^{(m+1)}(X_j^{(m+1)}) - q^{-(\beta_m,\beta_l+\beta_j)}\delta_l^{(m+1)}(X_j^{(m+1)})X_m^{(m+1)}]$
  $- q^{(\beta_l,\beta_m)}\delta_m^{(m+1)}(X_l^{(m+1)})X_j^{(m+1)}$.
  Since $\delta_l^{(m+1)}(X_j^{(m+1)})$ is homogeneous of degree $\beta_l + \beta_j$, we have
  $X_m^{(m+1)}\delta_l^{(m+1)}(X_j^{(m+1)}) - q^{-(\beta_m,\beta_l+\beta_j)}\delta_l^{(m+1)}(X_j^{(m+1)})X_m^{(m+1)} = \delta_m^{(m+1)} \circ \delta_l^{(m+1)}(X_j^{(m+1)})$.
  So, we get:

$$\begin{gathered} A + B - C = \\ (q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)})q^{-(\beta_l,\beta_j)} X_m^{(m+1)} X_j^{(m+1)} X_l^{(m+1)} \\ + q^{(\beta_l,\beta_m)} \delta_m^{(m+1)} \circ \delta_l^{(m+1)}(X_j^{(m+1)}) - q^{(\beta_l,\beta_m)} \delta_m^{(m+1)}(X_l^{(m+1)}) X_j^{(m+1)}. \end{gathered}$$

- Using the computation of $\delta_l^{(m+1)}(X_m^{(m+1)})$ made in the first point over, we can write:
  $D = q^{-[(\beta_m,\beta_j)+(\beta_l,\beta_j)]} X_j^{(m+1)} \delta_l^{(m+1)}(X_m^{(m+1)}) =$
  $q^{-(\beta_j,\beta_l+\beta_m)} X_j^{(m+1)}[(q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)}) X_m^{(m+1)} X_l^{(m+1)} - q^{(\beta_l,\beta_m)} \delta_m^{(m+1)}(X_l^{(m+1)})]$.
  So, we get:

  $$\begin{aligned} D = {} & q^{-(\beta_j,\beta_l+\beta_m)}(q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)}) X_j^{(m+1)} X_m^{(m+1)} X_l^{(m+1)} \\ & - q^{-(\beta_j,\beta_l+\beta_m)} q^{(\beta_l,\beta_m)} X_j^{(m+1)} \delta_m^{(m+1)}(X_l^{(m+1)}). \end{aligned}$$

- Now, we have
  $\delta_l^{(m+1)}(V) = A + B - C - D = q^{-(\beta_j,\beta_l}(q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)})[X_m^{(m+1)} X_j^{(m+1)} - q^{-(\beta_j,\beta_m)} X_j^{(m+1)} X_m^{(m+1)}] X_l^{(m+1)}$
  $+ q^{(\beta_l,\beta_m)}[q^{-(\beta_j,\beta_l+\beta_m)} X_j^{(m+1)} \delta_m^{(m+1)}(X_l^{(m+1)}) - \delta_m^{(m+1)}(X_l^{(m+1)}) X_j^{(m+1)}]$
  $+ q^{(\beta_l,\beta_m)} \delta_m^{(m+1)} \circ \delta_l^{(m+1)}(X_j^{(m+1)})$.
  So, we get:

  $$\begin{gathered} \delta_l^{(m+1)}(V) = q^{-(\beta_j,\beta_l)}(q^{(\beta_l,\beta_m)} - q^{-(\beta_l,\beta_m)}) V X_l^{(m+1)} \\ + q^{(\beta_l,\beta_m)}[q^{-(\beta_j,\beta_l+\beta_m)} X_j^{(m+1)} \delta_m^{(m+1)}(X_l^{(m+1)}) - \delta_m^{(m+1)}(X_l^{(m+1)}) X_j^{(m+1)}] \\ + q^{(\beta_l,\beta_m)} \delta_m^{(m+1)} \circ \delta_l^{(m+1)}(X_j^{(m+1)}) \end{gathered}$$

  As V is homogeneous of degree $\beta_j + \beta_m$, we have also

  $$\delta_l^{(m+1)}(V) = X_l^{(m+1)} V - q^{-(\beta_l,\beta_j+\beta_m)} V X_l^{(m+1)},$$

  which gives the required formula.

□

Now, let us observe that, if $j < l < m$, we have the following results:

- $\delta_m^{(m+1)}(X_l^{(m+1)}) \in k < X_{l+1}^{(m+1)}, \dots, X_{m-1}^{(m+1)} >$ (or is zero if $l = m-1$) (see section 3.2).
- Similarly, $V = \delta_m^{(m+1)}(X_j^{(m+1)}) \in k < X_{j+1}^{(m+1)}, \dots, X_{m-1}^{(m+1)} >$ (observe that $j < m-1$ by lemma 4.4. 1).
- $\delta_l^{(m+1)}(X_j^{(m+1)}) \in k < X_{j+1}^{(m+1)}, \dots, X_{l-1}^{(m+1)} >$ (or is zero if $l = j+1$), which implies that
  $\delta_m^{(m+1)} \circ \delta_l^{(m+1)}(X_j^{(m+1)}) \in k < X_{j+1}^{(m+1)}, \dots, X_{m-1}^{(m+1)} >$.

So, V, $\delta_m^{(m+1)}(X_l^{(m+1)})$, $\delta_l^{(m+1)} \circ \delta_l^{(m+1)}(X_j^{(m+1)})$, $X_l^{(m+1)}$, $X_j^{(m+1)}$ are all in $k < X_1^{(m+1)}, \dots, X_{m-1}^{(m+1)} >$ and, if we transform the equality of lemma 4.4. 2 by the algebra homomorphism $\Theta^{(m)}$, we obtain:

Lemma 4.4. 3

$$\begin{gathered} X_l^{(m)} U \ = \ q^{(\beta_l,\beta_m-\beta_j)} U X_l^{(m)} \\ + q^{(\beta_l,\beta_m)} \ [\ q^{-(\beta_j,\beta_l+\beta_m)} X_j^{(m)} \Theta^{(m)}(\delta_m^{(m+1)}(X_l^{(m+1)})) \ - \ \Theta^{(m)}(\delta_m^{(m+1)}(X_l^{(m+1)})) X_j^{(m)} \ ] \\ + q^{(\beta_l,\beta_m)} \Theta^{(m)}(\delta_m^{(m+1)} \circ \delta_l^{(m+1)}(X_j^{(m+1)})). \end{gathered}$$

- It results of the choice of j that $\Theta^{(m)}(\delta_m^{(m+1)}(X_l^{(m+1)})) \in \mathcal{P}^{(m)}$.

Assume that $l > j+1$, so that (by section 3.2)

$$\delta_l^{(m+1)}(X_j^{(m+1)}) \;=\; \sum_{\underline{a}\,=\,(a_{j+1},\ldots,a_{l-1})} c_{\underline{a}}(X_{j+1}^{(m+1)})^{a_{j+1}}\ldots(X_{l-1}^{(m+1)})^{a_{l-1}}$$

with each $c_{\underline{a}} \in k$.

As each $X_i^{(m+1)}$ is a h_m - eigenvector, each $\delta_m^{(m+1)}(X_{j+1}^{(m+1)})^{a_{j+1}}\ldots(X_{l-1}^{(m+1)})^{a_{l-1}})$ is a linear combination of products $M_1\delta_m^{(m+1)}(X_i^{(m+1)})M_2$ with $j < i < l$ and M_1 (resp. M_2) an ordered monomial in $X_{j+1}^{(m+1)}, \ldots , X_i^{(m+1)}$ (resp. in $X_i^{(m+1)}, \ldots , X_{l-1}^{(m+1)}$). This implies that $\Theta^{(m)}(\delta_m^{(m+1)} \circ \delta_l^{(m+1)}(X_j^{(m+1)}))$ is a linear combination of products $N_1\Theta^{(m)}(\delta_m^{(m+1)}(X_i^{(m+1)}))N_2$ with $j < i < l$, N_1 (resp. N_2) an ordered monomial in $X_{j+1}^{(m)}, \ldots , X_i^{(m)}$ (resp. in $X_i^{(m)}, \ldots , X_{l-1}^{(m)}$).
As $\Theta^{(m)}(\delta_m^{(m+1)}(X_i^{(m+1)})) \in \mathcal{P}^{(m)}$ for $j < i < m$, we conclude that

- $\Theta^{(m)}(\delta_m^{(m+1)} \circ \delta_l^{(m+1)}(X_j^{(m+1)})) \in \mathcal{P}^{(m)}$.

We observe that this result is also true when $l = j+1$ since, in this case, $\delta_l^{(m+1)}(X_j^{(m+1)})$ is zero (see section 3.2). So, if we transform the equality of lemma 4.4. 3 by the algebra homomorphism f_m, we obtain:

Lemma 4.4. 4

$$x_l^{(m)}u \;=\; q^{(\beta_l,\beta_m-\beta_j)}ux_l^{(m)}. \tag{16}$$

Recall (see section 3.2) that $V = \delta_m^{(m+1)}(X_j^{(m+1)}))$ is an homogeneous polynomial in $X_{j+1}^{(m+1)}, \ldots , X_{m-1}^{(m+1)}$ of degree $\beta_m + \beta_j$. So, since $\Theta^{(m)}$ transforms each $X_i^{(m+1)}$ $(1 \leq i < m)$ in $X_i^{(m)}$, $U = \Theta^{(m)}(V)$ is an homogeneous polynomial in $X_{j+1}^{(m)}, \ldots , X_{m-1}^{(m)}$ of degree $\beta_m + \beta_j$. This implies:

Lemma 4.4. 5
u is a nonzero polynomial in $x_{j+1}^{(m)}, \ldots , x_{m-1}^{(m)}$. Moreover, u is homogeneous of degree $\beta_m + \beta_j$.

Proof
$u = f_m(U)$ is nonzero by construction. Since f_m transforms each $X_i^{(m)}$ in $x_i^{(m)}$, it turns out that u is a polynomial in $x_{j+1}^{(m)}, \ldots , x_{m-1}^{(m)}$. Moreover, $u = f_m(U)$ is homogeneous of degree $\beta_m + \beta_j$ by lemma 4.3. 1

□

Recall (lemma 4.4. 1) that $j < l_p$ and that p is the greatest integer such that $l_p < m$. Denote by c the smallest integer such that $j < l_c$ $(1 \leq c \leq p)$. As $u \in k < x_{j+1}^{(m)}, \ldots , x_{m-1}^{(m)} >$, it results from (proposition 4.3. 2, 2.b.) that

$$u \;=\; u_1 z_{l_p}^{a_1} + \;\ldots\; + u_M z_{l_p}^{a_M}$$

with $M \geq 1$, $(a_1, \ldots , a_M) \in \mathbb{Z}^M$ and:
- If $p = 1$, then each $u_i \in k$.
- If $p > 1$ and $j \geq l_{p-1}$, then each $u_i \in k$.
- If $p > 1$ and $j < l_{p-1}$, then each $u_i \in k < x_{j+1}^{(l_p)}, \ldots , x_{l_{p-1}}^{(l_p)} >$,

so that we can write

- If $c = p$, then each $u_i \in k$.
- If $c < p$, then each $u_i \in k < x_{j+1}^{(l_p)}, \ldots , x_{l_{p-1}}^{(l_p)} >$.

In both cases, we may assume that $u_1, \ldots , u_M$ are all nonzero, that $a_1 < \ldots < a_M$, and we observe that

- Each $u_i \in k < x_1^{(l_p)},\ \dots\ ,\ x_{l_{p-1}}^{(l_p)} >$ and is homogeneous of degree $\beta_m + \beta_j - a_i \beta_{l_p}$.

In fact, since u is homogeneous of degree $\beta_m + \beta_j$ and $z_{l_p} = x_{l_p}^{(l_p)}$ is homogeneous of degree β_{l_p}, we have, for any $\rho \in \mathbb{Z}\Pi$,

$$\overline{h_\rho}(u) \ = \ q^{-(\rho,\ a_1\beta_{l_p})}\overline{h_\rho}(u_1) z_{l_p}^{a_1} + \ \dots\ + q^{-(\rho,\ a_M\beta_{l_p})}\overline{h_\rho}(u_M) z_{l_p}^{a_M}$$

$$= \ q^{-(\rho,\ \beta_m+\beta_j)}\ [u_1 z_{l_p}^{a_1} + \ \dots\ + u_M z_{l_p}^{a_M}]$$

and, by corollary 4.3. 2, we can identify the coefficients of $z_{l_p}^{a_i}$, so that $q^{-(\rho,\ a_i\beta_{l_p})}\overline{h_\rho}(u_i) = q^{-(\rho,\ \beta_m+\beta_j)} u_i$ and $\overline{h_\rho}(u_i) = q^{-(\rho,\ \beta_m+\beta_j - a_i\beta_{l_p})} u_i$.

By (proposition 4.3. 3, 1.) we have $x_{l_p}^{(m)} = z_{l_p} = x_{l_p}^{(l_p)}$. So, using lemma 4.4. 4 with $l = l_p$, we obtain:

$$z_{l_p} u \ = \ q^{(\beta_{l_p},\ \beta_m - \beta_j)} u z_{l_p}$$

and, by (lemma 4.3. 1, 4.),

$$z_{l_p} u \ = \ \overline{h_{l_p}}(u_1) z_{l_p}^{a_1+1} + \ \dots\ + \overline{h_{l_p}}(u_M) z_{l_p}^{a_M+1}.$$

This implies that

$$q^{(\beta_{l_p},\ \beta_m - \beta_j)}\ [u_1 z_{l_p}^{a_1} + \ \dots\ + u_M z_{l_p}^{a_M}] z_{l_p}$$

$$= \ [q^{-(\beta_{l_p},\ \beta_m+\beta_j-a_1\beta_{l_p})} u_1 z_{l_p}^{a_1} + \ \dots\ + q^{-(\beta_{l_p},\ \beta_m+\beta_j-a_N\beta_{l_p})} u_M z_{l_p}^{a_M}] z_{l_p}.$$

As over, if $1 \leq i \leq M$, we can identify the coefficients of $z_{l_p}^{a_i+1}$, so that, since $u_i \neq 0$ and q is not a root of unity, we have $-(\beta_{l_p}, \beta_m + \beta_j - a_i\beta_{l_p}) \ = \ (\beta_{l_p}, \beta_m - \beta_j)$.

So, $a_i \|\beta_{l_p}\|^2 = 2(\beta_{l_p}, \beta_m)$ and then $a_i \ = \ (\beta_{l_p}^\vee, \beta_m)$.

This implies that $M = 1$ and we conclude:

Lemma 4.4. 6

$$u \ = \ u_p z_{l_p}^{a_p}$$

with

- *If* $c = p$, *then* $u_p \ \in \ k^*$.
- *If* $c < p$, *then* $u_p \ \in \ k < x_{j+1}^{(l_p)},\ \dots\ ,\ x_{l_{p-1}}^{(l_p)} > \setminus \{0\}$.
- $a_p \ = \ (\beta_{l_p}^\vee, \beta_m)$.
- u_p *is homogeneous of degree* $\beta_m + \beta_j - a_p \beta_{l_p} = \gamma_p + \beta_j$ *with* $\gamma_p = s_{\beta_{l_p}}(\beta_m)$.

Now, we extend this result as follows:

Lemma 4.4. 7
Consider some integer d *with* $c \leq d \leq p$. *Then*

$$u \ = \ u_d z_{l_d}^{a_d} \dots z_{l_p}^{a_p}$$

with

- *If* $c = d$, *then* $u_d \ \in \ k^*$.
- *If* $c < d$, *then* $u_d \ \in \ k < x_{j+1}^{(l_d)},\ \dots\ ,\ x_{l_{d-1}}^{(l_d)} > \setminus \{0\}$.

- *Denote by* $(\gamma_d,\ \dots\ ,\ \gamma_p,\ \gamma_{p+1})$ *the sequence of (non necessarily positive) roots recursively defined by* $\gamma_{p+1} = \beta_m$ *and, for* $d \leq i \leq p$, $\gamma_i = s_{\beta_{l_i}}(\gamma_{i+1})$. *Then, each* $a_i = (\beta_{l_i}^\vee, \gamma_{i+1})$.
- u_d *is homogeneous of degree* $\gamma_d + \beta_j$.

Proof
We proceed by decreasing induction on d.
If $d = p$, everything has been done in lemma 4.4. 6.
Assume that $c \leq d < p$ and that the lemma is true at the rank $d + 1$, namely:

$$u \ = \ u_{d+1} z_{l_{d+1}}^{a_{d+1}} \ \dots \ z_{l_p}^{a_p}$$

with

- $u_{d+1} \ \in \ k < x_{j+1}^{(l_{d+1})},\ \dots\ ,\ x_{l_d}^{(l_{d+1})} > \setminus \{0\}$.
- If $(\gamma_{d+1},\ \dots\ ,\ \gamma_p,\ \gamma_{p+1})$ is the sequence of (non necessarily positive) roots recursively defined by $\gamma_{p+1} = \beta_m$ and, for $d + 1 \leq i \leq p$, $\gamma_i = s_{\beta_{l_i}}(\gamma_{i+1})$, then, each $a_i = (\beta_{l_i}^\vee, \gamma_{i+1})$.
- u_{d+1} is homogeneous of degree $\gamma_{d+1} + \beta_j$.

By (proposition 4.3. 2, 2.b.), we have:

$$u_{d+1} \ = \ v_1 z_{l_d}^{b_1} \ + \ \dots \ + \ v_M z_{l_d}^{b_M}$$

with $M \geq 1$, $(b_1,\ \dots\ ,\ b_M) \in \mathbb{Z}^M$ and:

- If $d = c$, then each $v_i \ \in \ k$.
- If $d > c$, then each $v_i \ \in \ k < x_{j+1}^{(l_d)},\ \dots\ ,\ x_{l_{d-1}}^{(l_d)} >$.

As over, we may assume that $v_1,\ \dots\ ,v_M$ are all nonzero, that $b_1 < \dots < b_M$, and observe that

- Each v_i is homogeneous of degree $\gamma_{d+1} + \beta_j - b_i \beta_{l_d}$.

By (proposition 4.3. 3, 2.), we have

$$x_{l_d}^{(m)} \ = \ z_{l_d} \ + \sum_{\underline{s} \ = \ (s_{d+1},\ \dots\ ,\ s_p) \ \in \ F} \eta(\underline{s}) z_{l_{d+1}}^{s_{d+1}} \ \dots \ z_{l_p}^{s_p},$$

where

- F is a finite (possibly empty) subset of $\mathbb{Z}^{p-d}$.
- If $\preceq$ denotes the inverse lexicographic order in $\mathbb{Z}^{p-d}$, then, for any $\underline{s}$ in F, we have $\underline{s} \ \prec \ 0$.
- For each $\underline{s}$ in F, we have $\eta(\underline{s}) \ \in \ k^*$.

By (lemma 4.3. 1, 4.), we have

$$z_{l_d} u_{d+1} \ = \ \overline{h_{l_d}}(v_1) z_{l_d}^{b_1 + 1} + \ \dots \ + \overline{h_{l_d}}(v_M) z_{l_d}^{b_M + 1}.$$

For any $\underline{e} \ = \ (e_{d+1},\ \dots\ ,\ e_p) \in \mathbb{Z}^{\,p-d}$, let us denote $z^{\underline{e}} := z_{l_{d+1}}^{e_{d+1}} \ \dots \ z_{l_p}^{e_p}$.
Again by (lemma 4.3. 1, 4.), there exists $(\lambda_{\underline{e},1},\ \dots\ ,\ \lambda_{\underline{e},M}) \in (k^*)^M$ such that

$$z^{\underline{e}} u_{d+1} \ = \ \lambda_{\underline{e},1} v_1 z_{l_d}^{b_1} z^{\underline{e}} + \ \dots \ + \lambda_{\underline{e},M} v_M z_{l_d}^{b_M} z^{\underline{e}}.$$

Now, if we set $\underline{a} \ = \ (a_{d+1},\ \dots\ ,\ a_p)$, we have

$$x^{(m)}_{l_d} u \ = \ (1) \ + \ (2)$$

with

$$(1) \ = \ z_{l_d} u_{d+1} z^{\underline{a}} \ = \ \overline{h_{l_d}}(v_1) z_{l_d}^{b_1+1} z^{\underline{a}} + \ \dots \ + \overline{h_{l_d}}(v_M) z_{l_d}^{b_M+1} z^{\underline{a}}$$

$$= \ q^{-(\beta_{l_d}, \ \gamma_{d+1}+\beta_j - b_1\beta_{l_d})} v_1 z_{l_d}^{b_1+1} z^{\underline{a}} + \ \dots \ + q^{-(\beta_{l_d}, \ \gamma_{d+1}+\beta_j - b_M\beta_{l_d})} v_M z_{l_d}^{b_M+1} z^{\underline{a}}$$

and

$$(2) \ = \ (\sum_{\underline{s} \ \in \ F} \eta(\underline{s}) z^{\underline{s}} \ ) \ u_{d+1} z^{\underline{a}} \ = \ \sum_{\underline{s} \ \in \ F} \eta(\underline{s}) \ (\lambda_{\underline{s},1} v_1 z_{l_d}^{b_1} z^{\underline{s}} + \ \dots \ + \lambda_{\underline{s},M} v_M z_{l_d}^{b_M} z^{\underline{s}} \ ) \ z^{\underline{a}}.$$

Besides, we also have

$$u x^{(m)}_{l_d} \ = \ (1') \ + \ (2')$$

with

$$(1') \ = \ u_{d+1} z^{\underline{a}} z_{l_d} \ = \ v_1 z_{l_d}^{b_1} z^{\underline{a}} z_{l_d} + \ \dots \ + v_M z_{l_d}^{b_M} z^{\underline{a}} z_{l_d}$$

$$= \ q^{-(a_{d+1}\beta_{l_{d+1}} \ + \ \dots \ + \ a_p\beta_{l_p}, \ \beta_{l_d})} (v_1 z_{l_d}^{b_1+1} + \ \dots \ + v_M z_{l_d}^{b_M+1}) z^{\underline{a}}$$

and

$$(2') \ = \ u_{d+1} z^{\underline{a}} \ [\sum_{\underline{s} \ \in \ F} \eta(\underline{s}) z^{\underline{s}} \ ] \ = \ \sum_{\underline{s} \ \in \ F} \eta(\underline{s}) (v_1 z_{l_d}^{b_1} z^{\underline{a}} z^{\underline{s}} + \ \dots \ + v_M z_{l_d}^{b_M} z^{\underline{a}} z^{\underline{s}}).$$

Using lemma 4.4. 4 and with $l = l_d$, we obtain:

$$(1) \ + \ (2) \ = \ q^{(\beta_{l_d}, \ \beta_m - \beta_j)} \ [\ (1') \ + \ (2') \ ].$$

We observe that (1) and $(1')$ are left linear combinations of monomials of type $z_{l_d}^h z^{\underline{a}}$ $(h \in \mathbb{Z})$ with coefficients in $A = k < x^{(l_d)}_{j+1}, \ \dots \ , x^{(l_d)}_{l_{d-1}} >$, while (2) and $(2')$ are left linear combinations of monomials of type $z_{l_d}^h z^{\underline{e}}$ $(h \in \mathbb{Z}, \ \underline{e} = \underline{a} + \underline{s}, \ s \in F)$ with coefficients in A. Recall that, for any $\underline{s} \in F$, we have $\underline{s} \prec 0 \ \Rightarrow \ \underline{e} = \underline{a} + \underline{s} \prec \underline{a}$. So, by corollary 4.3. 2, we have $(1) = q^{(\beta_{l_d}, \ \beta_m - \beta_j)}(1')$, which implies that, if $1 \le i \le M$,

$$q^{-(\beta_{l_d}, \ \gamma_{d+1}+\beta_j - b_i\beta_{l_d})} \ = \ q^{(\beta_{l_d}, \ \beta_m - \beta_j)} q^{-(a_{d+1}\beta_{l_{d+1}} \ + \ \dots \ + \ a_p\beta_{l_p}, \ \beta_{l_d})}.$$

Recall that, for $d+1 \le h \le p$, $\gamma_h = s_{\beta_{l_h}}(\gamma_{h+1}) = \gamma_{h+1} - (\beta^\vee_{l_h}, \gamma_{h+1})\beta_{l_h} = \gamma_{h+1} - a_h\beta_{l_h}$. If we add up all these equalities, we obtain $\gamma_{d+1} = \gamma_{p+1} - (a_{d+1}\beta_{l_{d+1}} \ + \ \dots \ + \ a_p\beta_{l_p})$. As $\gamma_{p+1} = \beta_m$, we get the formula

$$-(\beta_{l_d}, \ \gamma_{d+1} + \beta_j - b_i\beta_{l_d}) \ = \ (\beta_{l_d}, \ \beta_m - \beta_j) \ - \ (\beta_m \ - \ \gamma_{d+1}, \ \beta_{l_d})$$

for any $i \in [\![1, \ \dots \ , M]\!]$.

This implies that $b_i \|\beta_{l_d}\|^2 - (\beta_{l_d}, \ \gamma_{d+1}) = (\beta_{l_d}, \ \gamma_{d+1})$. So, as the integers b_i are distincts, we necessarily have $M = 1$, $b_1 = (\beta^\vee_{l_d}, \ \gamma_{d+1})$ and v_1 is homogeneous of degree $\gamma_{d+1} - b_1\beta_{l_d} + \beta_j = \gamma_d + \beta_j$ if we set $\gamma_d = \gamma_{d+1} - b_1\beta_{l_d} = s_{\beta_{l_d}}(\gamma_{d+1})$.

$\square$

Now, in the case $d = c$, lemma 4.4. 7 becomes:

$$u \ = \ u_c z_{l_c}^{a_c} \ \dots \ z_{l_p}^{a_p}$$

with

- $u_c \ \in \ k^*$.

- If $(\gamma_c, \ldots , \gamma_p, \gamma_{p+1})$ is the sequence of (non necessarily positive) roots recursively defined by $\gamma_{p+1} = \beta_m$ and, for $c \leq i \leq p$, $\gamma_i = s_{\beta_{l_i}}(\gamma_{i+1})$, then each $a_i = (\beta_{l_i}^\vee, \gamma_{i+1})$.

In this case, u_c is homogeneous of degree 0 and, since each $z_{l_d} = x_{l_d}^{(l_d)}$ is homogeneous of degree β_{l_d} (see lemma 4.3. 1), we obtain that u is also homogeneous of degree $a_c\beta_{l_c} + \ldots + a_p\beta_{l_p}$. So, as the degree of u is uniquely defined, we have

$$\beta_m + \beta_j = a_c\beta_{l_c} + \ldots + a_p\beta_{l_p}$$

and, by proposition 2.3. 3, we conclude:

Proposition 4.4. 1
If $\mathcal{P}^{(m)}$ *is not in* $Im(\phi_m)$, *then*

$$\Delta = \{i \in [\![1, \ldots , t]\!] \mid Z_i \in \mathcal{P}^{(2)}\}$$

is not a positive diagram with respect to (1) *in the sense of definition* 2.2. 2.

5 Connections between admissible and positive diagrams.

w and the algebra $R = U^+[w]$ are defined as in section 2. The admissible diagrams (with respect to the reduced expression (1)) are defined as in section 3.3.

We know that the longest element w_0 has a reduced expression of the form

$$w_0 = s_{\alpha_1} \ldots s_{\alpha_N} \qquad (\alpha_i \in \Pi \;\; for \;\; 1 \leq i \leq N = |\Phi^+|) \tag{17}$$

such that $s_{\alpha_1} \ldots s_{\alpha_t}$ is the reduced expression (1) of w.

In this section, we shall use the following conventions:

- Denote by $R_0 = U^+[w_0]$ the algebra associated to the reduced expression (17) as explained in section 2.1. and set $F_0 = Fract(R_0)$ it's division ring of fractions.
- As in section 3.1, we set $X_i = X_{\beta_i}$ for $1 \leq i \leq N$ and we observe that $X_1, \ldots , X_t$ are also the generators of $R = U^+[w]$ defined in section 2.1, so that $R = k < X_1, \ldots , X_t >$ is a subalgebra of $R_0 = k < X_1, \ldots , X_N >$.
- For any $\rho \in \mathbb{Z}\Pi$ we still denote by h_ρ the automorphism of R_0 defined as in section 3.1.
- Each h_ρ can uniquely be extended in an automorphism, still denoted h_ρ, of F_0.
- We still denote by H the set of all the automorphisms h_ρ of F_0 $(\rho \in \mathbb{Z}\Pi)$. It is a subgroup of $Aut(F_0)$.
- For any m in $[\![1, \ldots , N+1]\!]$, we define the algebra $R_0^{(m)}$ and it's canonical generators $X_1^{(m)}, \ldots , X_N^{(m)}$ as in section 3.2.

5.1 A property of admissible diagrams.

Consider Δ a diagram with respect to (1) (ie. a subset of $[\![1, \ldots , t]\!]$). It is also a diagram with respect to (17) but it is not quite clear whether the properties "Δ is admissible with respect to (1)" and "Δ is admissible with respect to (17)" are equivalent. In this section, we clarify this point.

Lemma 5.1. 1

1. *There exists a unique algebra isomorphism* g: $R = k < X_1, \ldots , X_t > \;\rightarrow\; k < X_1^{(t+1)}, \ldots , X_t^{(t+1)} > \;\subset R_0^{(t+1)}$ *which transforms each* X_i $(1 \leq i \leq t)$ *in* $X_i^{(t+1)}$.
2. *For any* $\rho \in \mathbb{Z}\Pi$ *and* $B \in R$, *we have* $g \circ h_\rho(B) = h_\rho \circ g(B)$.
3. *The map* $\mathcal{P} \mapsto g(\mathcal{P})$ *is a bijection from the set of* H *- invariant prime ideals of* R *onto the set of* H *- invariant prime ideals of* $k < X_1^{(t+1)}, \ldots , X_t^{(t+1)} >$.

Proof

1. We know (see section 3.1) that R is the k - algebra generated by the variables $X_1, \dots, X_t$ submitted to the Levendorskii - Soibelman relations (10) and (11). As $X_1^{(t+1)}, \dots, X_t^{(t+1)}$ satisfy the "same" relations (12) and (14) (see section 3.2), there exists a unique homomorphism g: $R = k < X_1, \dots, X_t > \ \rightarrow\ k < X_1^{(t+1)}, \dots, X_t^{(t+1)} > \ \subset\ R_0^{(t+1)}$ which transforms each X_i $(1 \leq i \leq t)$ in $X_i^{(t+1)}$. As g transforms the base of the ordered monomials in $X_1, \dots, X_t$ in the base of the ordered monomials in $X_1^{(t+1)}, \dots, X_t^{(t+1)}$, it is an isomorphism.
2. For $1 \leq i \leq t$, we have $g \circ h_\rho\ (X_i) = q^{-(\rho,\beta_i)} g\ (X_i) = q^{-(\rho,\beta_i)}\ X_i^{(t+1)} = h_\rho\ (X_i^{(t+1)}) = h_\rho \circ g\ (X_i)$. This implies that the homomorphisms $g \circ h_\rho$ and $h_\rho \circ g$ coincide on R.
3. results thoroughly of 1. and 2.

□

Until the end of this section, we identify the canonical generators $X_1, \dots, X_t$ of R with $X_1^{(t+1)}, \dots, X_t^{(t+1)}$ respectively, so that R is identified with the subalgebra $k < X_1^{(t+1)}, \dots, X_t^{(t+1)} >$ of $R_0^{(t+1)}$, H - *Spec*(R) is identified with the set of H - invariant prime ideals of $k < X_1^{(t+1)}, \dots, X_t^{(t+1)} >$ and the division ring $F = Fract(R)$ is identified with the subalgebra of F_0 defined as follows: $F = \{as^{-1}\ | a \in R,\ s \in R \setminus \{0\}\}$. ($R$ and F are both H - invariant.)

Lemma 5.1. 2
Consider an integer m *with* $2 \leq m \leq t+1$.
If $X_1^{(m)}, \dots, X_N^{(m)}$ *are the canonical generators of* $R_0^{(m)}$, *then* $X_1^{(m)}, \dots, X_t^{(m)}$ *are the canonical generators of* $R^{(m)}$.

Proof
We proceed by decreasing induction on m.

- If $m = t+1$, $X_1^{(t+1)}, \dots, X_t^{(t+1)}$ are the canonical generators of R by the previous identification. So, they also are the canonical generators of $R^{(t+1)}$ by the definition of this algebra.
- Assume that $2 \leq m \leq t$ and that $X_1^{(m+1)}, \dots, X_t^{(m+1)}$ are the canonical generators of $R^{(m+1)}$. Denote (temporarily) by $X_1'^{(m)}, \dots, X_t'^{(m)}$ the canonical generators of $R^{(m)}$. Recall (see section 3.2) that they are defined as follows:
- For each $i \in [\![1, \dots, t]\!]$, we have
  1. $m \leq i\ \Rightarrow\ X_i'^{(m)} = X_i^{(m+1)}$.
  2. $$i < m\ \Rightarrow\ X_i'^{(m)} = X_i^{(m+1)} + \sum_{l=1}^{+\infty} C_l^{(m+1)} (X_m^{(m+1)})^{-l}$$

     with
     $$C_l^{(m+1)} = \frac{(1-q_m)^{-l}}{[l]!_{q_m}}\ \lambda_{m,i}^{-l}\ (\delta_m^{(m+1)})^l\ (X_i^{(m+1)}).$$

Moreover, if h_m is the (unique) automorphism of $k < X_1^{(m+1)}, \dots, X_{m-1}^{(m+1)} >$ which satisfies $h_m(X_i^{(m+1)}) = \lambda_{m,i} X_i^{(m+1)}$ $(\lambda_{m,i} = q^{-(\beta_m,\beta_i)})$, we have $\delta_m^{(m+1)}(a) = X_m^{(m+1)} a - h_m(a) X_m^{(m+1)}$ for any $a \in k < X_1^{(m+1)}, \dots, X_{m-1}^{(m+1)} >$. As $X_1^{(m)}, \dots, X_t^{(m)}$ are defined by the same formulas, the proof is over.

□

Recall (see sections 3.1 and 3.2) that each prime ideal of $R_0^{(t+1)}$ (resp. R) is completely prime. So, if $\mathcal{Q}$ is any prime ideal of $R_0^{(t+1)}$, then $\mathcal{P} = \mathcal{Q} \cap R$ is a prime ideal of R. Moreover, since $R = k < X_1^{(t+1)}, \dots, X_t^{(t+1)} >$ is H - invariant, we have: ($\mathcal{Q}$ is H - invariant) $\Rightarrow$ ($\mathcal{P}$ is H - invariant).

Lemma 5.1. 3
If $\mathcal{P}$ *is any* H *- invariant prime ideal of* R, *there exists a* H *- invariant prime ideal* $\mathcal{Q}$ *of* $R_0^{(t+1)}$, *such that*

- $\mathcal{P} = \mathcal{Q} \cap R$.
- $\mathcal{Q} \cap \{X_{t+1}^{(t+1)}, \ \dots \ , \ X_N^{(t+1)}\} = \emptyset$.

Proof
Set

$$\mathcal{Q} \ = \sum_{\underline{a} \ = \ (a_{t+1}, \ \dots \ , \ a_N)} \mathcal{P}(X^{(t+1)})^{\underline{a}}$$

with $\underline{a} \in \mathbb{N}^{N-t}$ and $(X^{(t+1)})^{\underline{a}} := (X_{t+1}^{(t+1)})^{a_{t+1}} \ \dots \ (X_N^{(t+1)})^{a_N}$.

Observe that, by the results recalled in 3.2, the family $((X^{(t+1)})^{\underline{a}})$ $(\underline{a} = (a_{t+1}, \dots , a_N) \in \mathbb{N}^{N-t})$ is a basis of $R_0^{(t+1)}$ as a right R - module. This implies that $\mathcal{Q} \cap R = \mathcal{P}$ and, since $1 \notin \mathcal{P}$, that each $(X^{(t+1)})^{\underline{a}} \notin \mathcal{Q}$ $(\underline{a} \in \mathbb{N}^{N-t})$. In particular, $\mathcal{Q} \cap \{X_{t+1}^{(t+1)}, \ \dots \ , \ X_N^{(t+1)}\} = \emptyset$.
For each $\underline{a} = (a_{t+1}, \dots , a_N) \in \mathbb{N}^{N-t}$ and each $u \in R$, we have $(X^{(t+1)})^{\underline{a}}\, u = h_{\underline{a}}(u)\ (X^{(t+1)})^{\underline{a}}$ with $h_{\underline{a}} = h_{a_{t+1}} \circ \dots \circ h_{a_N} \in H$ (see section 3.2).
This implies that $\mathcal{Q}$ is an R - bimodule.

Since, for any $\underline{a}$ and $\underline{b}$ in $\mathbb{N}^{N-t}$, we have $(X^{(t+1)})^{\underline{a}}\ (X^{(t+1)})^{\underline{b}} = \lambda\ (X^{(t+1)})^{\underline{c}}$ with $\underline{c} = \underline{a} + \underline{b}$ and $\lambda \in k^*$, $\mathcal{Q}$ is a right $R_0^{(t+1)}$ - module. Moreover, since $(X^{(t+1)})^{\underline{a}}\ \mathcal{P} = h_{\underline{a}}(\mathcal{P})\ (X^{(t+1)})^{\underline{a}} = \mathcal{P}\ (X^{(t+1)})^{\underline{a}}$, we obtain that $\mathcal{Q}$ is an ideal in $R_0^{(t+1)}$.

Now, consider A, B in $R_0^{(t+1)}$ and assume that $AB \in \mathcal{Q}$. As in section 4.3, denote by $\preceq$ the inverse lexicographic order in $\mathbb{N}^{N-t}$ and assume that, neither A, nor B is in $\mathcal{Q}$. Write

$$A \ = \sum_{\underline{a} \ \in F} A_{\underline{a}}(X^{(t+1)})^{\underline{a}}$$

with F a non empty finite subset of $\mathbb{N}^{N-t}$ and each $A_{\underline{a}}$ in R. Since, $A \notin \mathcal{Q}$, at least one coefficient $A_{\underline{a}}$ is not in $\mathcal{P}$. If we choose such a coefficient with $\underline{a}$ minimal, we get:

$$A \ = \ A_1 \ + \ A_{\underline{a}}(X^{(t+1)})^{\underline{a}} \ + \sum_{\underline{a'} \ \in F'} A_{\underline{a'}}(X^{(t+1)})^{\underline{a'}}$$

with $A_1 \in \mathcal{Q}$ and $\underline{a} \prec \underline{a'}$ for any $\underline{a'}$ in F'. Similarly, we have

$$B \ = \ B_1 \ + \ B_{\underline{b}}(X^{(t+1)})^{\underline{b}} \ + \sum_{\underline{b'} \ \in G'} B_{\underline{b'}}(X^{(t+1)})^{\underline{b'}}$$

with $B_1 \in \mathcal{Q}$, $B_{\underline{b}}$ is not in $\mathcal{P}$ and $\underline{b} \prec \underline{b'}$ for any $\underline{b'}$ in G'.

Now, we have

$$(A \ - \ A_1)(B \ - \ B_1) \ = \ \lambda A_{\underline{a}} B_{\underline{b}}(X^{(t+1)})^{\underline{c}} \ + \sum_{\underline{c'} \ \in E'} C_{\underline{c'}}(X^{(t+1)})^{\underline{c'}}$$

with λ in k^*, $\underline{c} = \underline{a} + \underline{b}$, each $C_{\underline{c'}}$ in R and, for each $\underline{c'}$ in E', $\underline{c} \prec \underline{c'}$. As $(A \ - \ A_1)(B \ - \ B_1) \in \mathcal{Q}$, this implies that $A_{\underline{a}}B_{\underline{b}} \in \mathcal{P}$, which is impossible since $\mathcal{P}$ is completely prime. This implies that $\mathcal{Q}$ is (completely) prime.

Since $\mathcal{P}$ is H - invariant and each monomial $X^{(t+1)})^{\underline{a}}$ $(\underline{a} \in \mathbb{N}^{N-t})$, is a H - eigenvector, $\mathcal{Q}$ is H - invariant.

□

Lemma 5.1. 4
Consider $\mathcal{P}_0 \in Spec(R_0)$ *and* $\mathcal{P} \in Spec(R)$.
Assume that $\mathcal{P} = \mathcal{Q} \cap R$ *with* $\mathcal{Q} = \mathcal{P}_0^{(t+1)}$ *= the canonical image of* $\mathcal{P}_0$ *in* $Spec(R_0^{(t+1)})$.
For any $m \in [\![2, \dots , t+1]\!]$, *we denote by* $\mathcal{P}^{(m)}$ *(resp.* $\mathcal{P}_0^{(m)}$*) the canonical image of* $\mathcal{P}$ *(resp.* $\mathcal{P}_0$*) in* $Spec(R^{(m)})$ *(resp.* $Spec(R_0^{(m)})$*) and we recall that, by lemma 5.1. 2,* $R^{(m)}$ *is the subalgebra of* $R_0^{(m)}$ *generated by it's* t *first canonical generators* $X_1^{(m)}, \ \dots \ , \ X_t^{(m)}$. *Then*

$$\mathcal{P}^{(m)} \;=\; \mathcal{P}_0^{(m)} \;\cap\; R^{(m)}.$$

Proof
We prove this by decreasing induction on m.

- If $m = t+1$, we have $R^{(m)} = R$, $\mathcal{P}^{(m)} = \mathcal{P}$ and $\mathcal{P}_0^{(m)} = \mathcal{Q}$. So, we have the required equality by assumption.
- Assume $m \leq t$ and $\mathcal{P}^{(m+1)} \;=\; \mathcal{P}_0^{(m+1)} \;\cap\; R^{(m+1)}$.
  - Assume that $X_m^{(m+1)} \notin \mathcal{P}_0^{(m+1)}$, so that $X_m^{(m+1)} \notin \mathcal{P}^{(m+1)}$.
    Set $S_m = \{(X_m^{(m+1)})^d \mid d \in \mathbb{N}\}$ and recall (sections 3.2 and 3.3) that
    - $R^{(m)}\, S_m^{-1} = R^{(m+1)}\, S_m^{-1}$.
    - $\mathcal{P}^{(m)} = \mathcal{P}^{(m+1)}\, S_m^{-1} \;\cap\; R^{(m)}$.
    - $\mathcal{P}_0^{(m)} = \mathcal{P}_0^{(m+1)}\, S_m^{-1} \;\cap\; R_0^{(m)}$.

    As $R^{(m)} \;\subseteq\; R_0^{(m)}$ and $\mathcal{P}^{(m+1)} \;\subseteq\; \mathcal{P}_0^{(m+1)}$, we have

    $$\mathcal{P}^{(m)} \;=\; \mathcal{P}^{(m+1)} S_m^{-1} \;\cap\; R^{(m)} \;\subseteq\; \mathcal{P}_0^{(m+1)} S_m^{-1} \;\cap\; R_0^{(m)} \;=\; \mathcal{P}_0^{(m)}.$$

    This implies that

    $$\mathcal{P}^{(m)} \;\subseteq\; \mathcal{P}_0^{(m)} \;\cap\; R^{(m)}.$$

    Now, consider some $u \in \mathcal{P}_0^{(m)} \;\cap\; R^{(m)}$.
    We have $\mathcal{P}_0^{(m)} \;\subseteq \mathcal{P}_0^{(m+1)}\, S_m^{-1} \;\Rightarrow\; u\,(X_m^{(m+1)})^{d_1} \in \mathcal{P}_0^{(m+1)}$ with $d_1 \in \mathbb{N}$.
    We have $R^{(m)} \;\subseteq R^{(m+1)}\, S_m^{-1} \;\Rightarrow\; u\,(X_m^{(m+1)})^{d_2} \in R^{(m+1)}$ with $d_2 \in \mathbb{N}$.
    Since $m \leq t$, $X_m^{(m+1)}$ is in $R^{(m+1)}$ and so, with $d = \max\,(d_1, d_2)$, we obtain that $u\,(X_m^{(m+1)})^d \in \mathcal{P}_0^{(m+1)} \;\cap\; R^{(m+1)} = \mathcal{P}^{(m+1)} \;\Rightarrow\; u \in \mathcal{P}^{(m+1)}\, S_m^{-1} \;\cap\; R^{(m)} = \mathcal{P}^{(m)}$. This implies that

    $$\mathcal{P}^{(m)} \;=\; \mathcal{P}_0^{(m)} \;\cap\; R^{(m)}.$$

  - Assume that $X_m^{(m+1)} \in \mathcal{P}_0^{(m+1)}$, so that, since $m \leq t$,

    $$X_m^{(m+1)} \;\in\; \mathcal{P}_0^{(m+1)} \;\cap\; R^{(m+1)} \;=\; \mathcal{P}^{(m+1)}.$$

    Denote by

    $$g:\; R^{(m)} \;\to\; \frac{R^{(m+1)}}{\mathcal{P}^{(m+1)}}$$

    the only algebra homomorphism which transforms each $X_i^{(m)}$ $(1 \leq i \leq t)$ in $g(X_i^{(m)}) = X_i^{(m+1)} + \mathcal{P}^{(m+1)}$ (the canonical image of $X_i^{(m+1)}$ in $\dfrac{R^{(m+1)}}{\mathcal{P}^{(m+1)}}$). We recall (section 3.3) that

    $$\mathcal{P}^{(m)} \;=\; ker(g).$$

    Denote by

    $$g_0:\; R_0^{(m)} \;\to\; \frac{R_0^{(m+1)}}{\mathcal{P}_0^{(m+1)}}$$

the only algebra homomorphism which transforms each $X_i^{(m)}$ $(1 \leq i \leq N)$ in $g_0(X_i^{(m)}) = X_i^{(m+1)} + \mathcal{P}_0^{(m+1)}$ (the canonical image of $X_i^{(m+1)}$ in $\frac{R_0^{(m+1)}}{\mathcal{P}_0^{(m+1)}}$), so that

$$\mathcal{P}_0^{(m)} \; = \; ker(g_0).$$

The canonical homomorphism $R_0^{(m+1)} \; \rightarrow \; \frac{R_0^{(m+1)}}{\mathcal{P}_0^{(m+1)}}$ restricted to $R^{(m+1)}$ has it's kernel equal to $\mathcal{P}_0^{(m+1)} \cap R^{(m+1)} = \mathcal{P}^{(m+1)}$. So, it induces an injective homomorphism

$$\epsilon : \; \frac{R^{(m+1)}}{\mathcal{P}^{(m+1)}} \; \rightarrow \; \frac{R_0^{(m+1)}}{\mathcal{P}_0^{(m+1)}}$$

which transforms each $X_i^{(m+1)} + \mathcal{P}^{(m+1)}$ $(1 \leq i \leq t)$ in $X_i^{(m+1)} + \mathcal{P}_0^{(m+1)}$. So,

$$\epsilon \circ g : \; R^{(m)} \; \rightarrow \; \frac{R_0^{(m+1)}}{\mathcal{P}_0^{(m+1)}}$$

is an algebra homomorphism which transforms each $X_i^{(m)}$ $(1 \leq i \leq t)$ in $X_i^{(m+1)} + \mathcal{P}_0^{(m+1)} = g_0\ (X_i^{(m)})$. This implies that $\epsilon \circ g$ is the restriction of g_0 to $R^{(m)}$ and so, that $Ker\ (\epsilon \circ g) = \mathcal{P}_0^{(m)} \; \cap \; R^{(m)}$. Since ϵ is injective, we also have $Ker(\epsilon \circ g) = Ker\ (g) = \mathcal{P}^{(m)}$ and the proof is over.

□

Proposition 5.1. 1
Consider a diagram Δ *with respect to* (1) (*ie. a subset of* $[\![1, \; ... \; , \; t]\!]$), *so that* Δ *is also a diagram with respect to* (17). *Then* Δ *is admissible with respect to* (1) *if and only if* Δ *is admissible with respect to* (17).

Proof

- Assume that Δ is admissible with respect to (17). This means (see section 3.3) that there exists some $\mathcal{P}_0$ in $Spec(R_0)$ such that it's canonical image $\mathcal{P}_0^{(2)}$ in $Spec(R_0^{(2)})$ verifies:

  $$\mathcal{P}_0^{(2)} \; \cap \; \{Z_1, \; ... \; , \; Z_N\} \; = \; \{Z_i \mid i \; \in \; \Delta\}.$$

  Denote by $\mathcal{Q} = \mathcal{P}_0^{(t+1)}$ the canonical image of $\mathcal{P}_0$ in $Spec(R_0^{(t+1)})$ and set $\mathcal{P} = \mathcal{Q} \; \cap \; R$. We know that $\mathcal{P} \in Spec(R)$ and, by lemma 5.1. 4, it's canonical image $\mathcal{P}^{(2)}$ in $Spec(R^{(2)})$ verifies

  $$\mathcal{P}^{(2)} \; = \; \mathcal{P}_0^{(2)} \; \cap \; R^{(2)}.$$

  So,

  $$\mathcal{P}^{(2)} \; \cap \; \{Z_1, \; ... \; , \; Z_t\} \; = \; \mathcal{P}_0^{(2)} \; \cap \; R^{(2)} \; \cap \; \{Z_1, \; ... \; , \; Z_t\} \; = \; \mathcal{P}_0^{(2)} \; \cap \; \{Z_1, \; ... \; , \; Z_t\} \; .$$

  Since Δ is a subset of $[\![1, \; ... \; , \; t]\!]$), this implies that

  $$\mathcal{P}^{(2)} \; \cap \; \{Z_1, \; ... \; , \; Z_t\} \; = \; \{Z_i \mid i \; \in \; \Delta\}.$$

  So Δ is admissible with respect to (1).

- Assume that Δ is admissible with respect to (1). This implies (see section 3.3) that there exists some $\mathcal{P}$ in $H - Spec(R)$ whose canonical image $\mathcal{P}^{(2)}$ in $Spec(R^{(2)})$ is the ideal of $R^{(2)}$ generated by $\{Z_i \mid i \; \in \; \Delta\}$ $(\mathcal{P}^{(2)} = 0$ if $\Delta = \emptyset)$.
  Since $\mathcal{P}$ is $H -$ invariant, it results from lemma 5.1. 3 that there exists a ($H -$ invariant) prime ideal $\mathcal{Q}$ of $R_0^{(t+1)}$, such that

- $\mathcal{P} = \mathcal{Q} \cap R$.
- $\mathcal{Q} \cap \{X_{t+1}^{(t+1)}, \ldots, X_N^{(t+1)}\} = \emptyset$.

Now, by corollary 4.1. 1, there exists $\mathcal{P}_0$ in $Spec(R_0)$ whose canonical image $\mathcal{P}_0^{(t+1)}$ in $Spec(R_0^{(t+1)})$ is equal to $\mathcal{Q}$ and whose canonical image $\mathcal{P}_0^{(2)}$ in $Spec(R_0^{(2)})$ does not meet the set $\{Z_{t+1}, \ldots, Z_N\}$.
As over, by lemma 5.1. 4, we have

$$\mathcal{P}^{(2)} = \mathcal{P}_0^{(2)} \cap R^{(2)}.$$

Since $\mathcal{P}_0^{(2)}$ does not meet the set $\{Z_{t+1}, \ldots, Z_N\}$, we have

$$\mathcal{P}_0^{(2)} \cap \{Z_1, \ldots, Z_N\} = \mathcal{P}_0^{(2)} \cap \{Z_1, \ldots, Z_t\} .$$

So, since $\{Z_1, \ldots, Z_t\} \subseteq R^{(2)}$, we can write

$$\begin{aligned} \mathcal{P}_0^{(2)} \cap \{Z_1, \ldots, Z_N\} &= \mathcal{P}_0^{(2)} \cap R^{(2)} \cap \{Z_1, \ldots, Z_t\} \\ &= \mathcal{P}^{(2)} \cap \{Z_1, \ldots, Z_N\} = \{Z_i \mid i \in \Delta\}. \end{aligned}$$

This proves that Δ is admissible with respect to (17).

□

5.2 The case $w = w_0$.

Assume (in this subsection only) that $w = w_0$.

Lemma 5.2. 1
For any diagram Δ (*with respect to* (1)) *we have:*

$$\Delta \text{ positive} \Rightarrow \Delta \text{ admissible}.$$

Proof
As in section 3.3, we denote by $\mathcal{P}_\Delta^{(2)}$ the ideal of $\overline{R} = R^{(2)}$ generated by the canonical generators Z_i with $i \in \Delta$ ($\mathcal{P}_\Delta^{(2)} = 0$ if $\Delta = \emptyset$). Recall (see section 3.3) that $\mathcal{P}_\Delta^{(2)} \in H - Spec(R^{(2)})$.
Consider an integer m with $2 \leq m \leq t+1$. We first prove, by induction on m, that there exists a prime ideal ideal $\mathcal{P}^{(m)} \in Spec(R^{(m)})$ such that $\mathcal{P}_\Delta^{(2)}$ is the canonical image of $\mathcal{P}^{(m)}$ in $Spec(R^{(2)})$.

- If $m = 2$, we just have to choose $\mathcal{P}^{(2)} = \mathcal{P}_\Delta^{(2)}$.

- Assume that, for some m with $2 \leq m \leq t$, there exists a prime ideal ideal $\mathcal{P}^{(m)} \in Spec(R^{(m)})$ such that $\mathcal{P}_\Delta^{(2)}$ is the canonical image of $\mathcal{P}^{(m)}$ in $Spec(R^{(2)})$. Since Δ is positive, we deduce from proposition 4.4. 1 that $\mathcal{P}^{(m)} = \phi_m(\mathcal{P}^{(m+1)})$ with $\mathcal{P}^{(m+1)}$ in $Spec(R^{(m+1)})$.
  So, $\mathcal{P}_\Delta^{(2)}$ is the canonical image of $\mathcal{P}^{(m+1)}$ in $Spec(R^{(2)})$, and our affirmation is proved.
  In particular, there exists $\mathcal{P} \in Spec(R^{(t+1)}) = Spec(R)$ such that $\mathcal{P}_\Delta^{(2)}$ is the canonical image of $\mathcal{P}$ in $Spec(R^{(2)})$. This means (see section 3.3) that Δ is admissible.

□

Proposition 5.2. 1
The positive diagrams (*with respect to* (1)) *coincide with the admissible diagrams* (*with respect to* (1)).

Proof
By lemma 5.2 1, the set of positive diagrams is contained in the set of admissible diagrams. By proposition 2.3. 1, the number of positive diagrams is equal to the number of u in the Weyl group W such that $u \leq w$ (where $\leq$ denotes the Bruhat order). Since we assume that $w = w_0$, we get that the number of positive diagrams is equal to the cardinal $|W|$ of W. Since the set of admissible diagrams is in one to one correspondence with $H - Spec(R) = H - Spec(U^+)$, the number of admissible diagrams does not depend on the reduced decomposition of w. In [17], Antoine Mériaux gives a precise description of admissible diagrams, for each type of the simple complex Lie algebra $\mathfrak{g}$, and for a particular reduced decomposition of w_0 so that he can compute their number and check that it is precisely equal to $|W|$. This implies that the positive diagrams coincide with the admissible diagrams.

□

Let us observe that, in this proof, we use a result of Antoine Mériaux to prove that the number of admissible diagrams (which is also the cardinal of $H - Spec(U^+)$) is equal to $|W|$. This equality can also be obtained using results of M. Gorelik, N. Andruskiewitsch and F. Dumas ([8] and [2]) or T. J. Hodges and T. Levasseur and M. Toro [10] but this should require some (minor) restricted assumptions on the choice of the ground field k (char$(k) = 0$) or on the parameter q (q transcendent).

5.3 The general case.

We can now prove, in the general case (ie. w is not any longer assumed to be the longest element of W):

Theorem 5.3. 1
The positive diagrams (with respect to (1)) coincide with the admissible diagrams (with respect to (1)).

Proof
Consider a diagram Δ with respect to (1) (ie. a subset of $[\![1, \dots , t]\!]$), so that Δ is also a diagram with respect to (17).

- Assume that Δ is positive (with respect to (1)). By proposition 2.3. 2, Δ is positive with respect to (17) and, by proposition 5.2. 1, Δ is admissible with respect to (17). Now, by proposition 5.1. 1, Δ is admissible with respect to (1).
- Assume that Δ is admissible (with respect to (1)). By proposition 5.1. 1, Δ is admissible with respect to (17) and, by proposition 5.2. 1, Δ is positive with respect to (17). Now, by proposition 2.3. 2, Δ is positive with respect to (1).

□

Corollary 5.3. 1 (*All the diagrams are with respect to* (1).)

1. *The map* $\zeta: \quad \Delta = \{j_1, \dots , j_s\} \mapsto u = w^{\Delta} = s_{\alpha_{j_1}} \circ \dots \circ s_{\alpha_{j_s}}$ *is a bijection from the set of admissible diagrams onto the set* $\{u \in W \mid u \leq w\}$ $(\zeta(\emptyset) = Id)$.
2. *Consider an admissible diagram* $\Delta = \{j_1, \dots , j_s\}$ *and some integer* $i \in [\![1, \dots , t]\!]$. *Set* $\Delta \cap [\![i+1, \dots , t]\!] = \{j_c, \dots , j_s\}$ $(1 \leq c \leq s)$. *Then the expression* $s_{\alpha_i} \circ s_{\alpha_{j_c}} \circ \dots \circ s_{\alpha_{j_s}}$ *is reduced. In particular,* $s_{\alpha_{j_1}} \circ \dots \circ s_{\alpha_{j_s}}$ *is a reduced expression of* $\zeta(\Delta)$.
3. *Consider some* $u \in W$ *with* $u \leq w$. *Then, the only admissible diagram* Δ *such that* $\zeta(\Delta) = u$ *is recursively defined as follows:*
   -
     $$1 \in \Delta \Leftrightarrow l(s_{\alpha_1} \circ u) = l(u) - 1 \Leftrightarrow u^{-1}(\alpha_1) \text{ is a negative root.}$$
   - *Consider some integer* $i \in [\![1, \dots , t-1]\!]$ *and assume that* $\Delta \cap [\![1, \dots , i]\!] = \{j_1, \dots , j_d\}$. *Set* $u_i = s_{\alpha_{j_d}} \circ \dots \circ s_{\alpha_{j_2}} \circ s_{\alpha_{j_1}} \circ u$ ($u_i = u$ *if* $\Delta \cap [\![1, \dots , i]\!] = \emptyset$). *Then*

     $$i+1 \in \Delta \Leftrightarrow l(s_{\alpha_{i+1}} \circ u_i) = l(u_i) - 1 \Leftrightarrow u_i^{-1}(\alpha_{i+1}) \text{ is a negative root.}$$

Proof
As the admissible diagrams coincide with the positive diagrams, 1. is proposition 2.3. 1 (assertion 2.) and 2. is proposition 2.2. 2. Now, let us prove 3.

- Assume that $1 \in \Delta$, so that $\Delta = \{j_1, \dots , j_s\}$ with $j_1 = 1$. As $u = \zeta(\Delta) = s_{\alpha_{j_1}} \circ \dots \circ s_{\alpha_{j_s}}$, we have $l(s_{\alpha_1} \circ u) = l(s_{\alpha_{j_2}} \circ \dots \circ s_{\alpha_{j_s}}) = s - 1 = l(u) - 1$.
Conversely, assume that $l(s_{\alpha_1} \circ u) = l(u) - 1$. If $1 \notin \Delta$, then $\Delta \cap [\![2, \dots , t]\!] = \Delta$ and, by 2., $l(s_{\alpha_1} \circ u) = l(u) + 1$. So, we obtain a contradiction.
Now, we have classically: $l(s_{\alpha_1} \circ u) = l(u) - 1 \;\Leftrightarrow\; l(u^{-1} \circ s_{\alpha_1}) = l(u^{-1}) - 1 \;\Leftrightarrow\; u^{-1}(\alpha_1)$ is a negative root.

- Assume that $i + 1 \in \Delta$, so that $\Delta \cap [\![i + 1, \dots , t]\!] = \{j_{d+1}, \dots , j_s\}$ with $j_{d+1} = i + 1$. As $u_i = s_{\alpha_{j_d}} \circ \dots \circ s_{\alpha_{j_2}} \circ s_{\alpha_{j_1}} \circ u = s_{\alpha_{j_{d+1}}} \circ \dots \circ s_{\alpha_{j_s}}$, we have $l(s_{\alpha_{i+1}} \circ u) = l(s_{\alpha_{j_{d+2}}} \circ \dots \circ s_{\alpha_{j_s}}) = s - (d + 1) = l(u_i) - 1$.
Conversely, assume that $l(s_{\alpha_{i+1}} \circ u_i) = l(u_i) - 1$. If $i + 1 \notin \Delta$, then $\Delta \cap [\![i + 2, \dots , t]\!] = \{j_{d+1}, \dots , j_s\}$ and, by 2., $l(s_{\alpha_{i+1}} \circ u_i) = l(u_i) + 1$. So, we still have a contradiction and we conclude as over.

□

From corollary 5.3. 1 (assertions 1. and 2.) and lemma 2.2. 1 (assertion 2.), we deduce:

Corollary 5.3. 2

1. *The map* ζ' : $\Delta = \{j_1, \dots , j_s\} \;\mapsto\; u' = v^{\Delta} = s_{\alpha_{j_s}} \circ \dots \circ s_{\alpha_{j_1}}$ *is a bijection from the set of admissible diagrams onto the set* $\{u \in W \mid u \;\leq\; v \;=\; w^{-1}\}$ $(\zeta'(\Delta) = (\zeta(\Delta))^{-1},\;\; \zeta'(\emptyset) = Id)$.
2. *Consider an admissible diagram* $\Delta = \{j_1, \dots , j_s\}$ *and some integer* $i \in [\![1, \dots , t]\!]$. *Set* $\Delta \cap [\![i + 1, \dots , t]\!] = \{j_c, \dots , j_s\}$ $(1 \;\leq\; c \;\leq\; s)$. *Then the expression* $s_{\alpha_{j_s}} \circ \dots \circ s_{\alpha_{j_c}} \circ s_{\alpha_i}$ *is reduced. In particular,* $s_{\alpha_{j_s}} \circ \dots \circ s_{\alpha_{j_1}}$ *is a reduced expression of* $\zeta'(\Delta)$.

If w and it's reduced decomposition are chosen as in the example of section 2.1, we know (proposition 2.1. 1) that $U^+[w]$ is the quantum matrices algebra $O_q(M_{p,m}(k))$ with $m = n - p + 1$ and with the same canonical generators (as $U^+[w]$). So, by [5], the admissible diagrams are the ⅃ - diagrams. In this case, corollary 5.3. 2 has also been proved (with quite different methods) by A. Postnikov ([18], theorem 19.1.) and by T. Lam and L. Williams ([15], theorem 5.3.).

In the general case, using the bijection between the set of admissible diagrams and $H - Spec(U^+[w])$ recalled in section 3.3, the two bijections of proposition 2.3. 1 and the coincidence of admissible and positive diagrams (theorem 5.3. 1), we get

Corollary 5.3. 3
1. There exists a natural bijection from the set $\{u \in W \mid u \;\leq\; v \;=\; w^{-1}\}$ *onto* $H - Spec(U^+[w])$. *It is defined by*

$$v^{\Delta} \;\mapsto\; \mathcal{P}_{\Delta}$$

where Δ *describes the set of admissible diagrams.*
2. There exists a natural bijection from the set $\{u \in W \mid u \;\leq\; w\}$ *onto* $H - Spec(U^+[w])$. *It is defined by*

$$w^{\Delta} \;\mapsto\; \mathcal{P}_{\Delta}$$

where Δ *describes the set of admissible diagrams.*

When $U^+[w]$ is the quantum matrices algebra $O_q(M_{p,m}(k))$ as over, S. Launois [13] has constructed (with different methods) a bijection from $\{u \in W \mid u \;\leq\; v\}$ onto $H - Spec(O_q(M_{p,m}(k)))$ which, moreover, preserves the ordering (where the Weyl group is provided with the Bruhat order and $H - Spec(O_q(M_{p,m}(k)))$ is provided with the inclusion). So, it seems natural to ask the following questions:

Question 5.3. 1
Assume w *and it's reduced decomposition chosed as in the example of section 2.1.*
Does the S. Launois bijection from $\{u \in W \mid u \;\leq\; v\}$ *onto* $H - Spec(O_q(M_{p,m}(k))) \;=\; H - Spec(U^+[w])$ *coincide with the first bijection of corollary 5.3. 3 ? If not, is there a simple relation between those two bijections ?*

Question 5.3. 2
In the general case, are the two bijections of corollary 5.3. 3 isomorphisms of ordered sets, where the set $\{u \in W \mid u \;\leq\; v\}$ *(resp.* $\{u \in W \mid u \;\leq\; w\}$*) is provided with the Bruhat order and* $H - Spec(U^+[w])$ *is provided with the inclusion?*

References

[1] J. Alev et F. Dumas, *Sur le corps des fractions de certaines algèbres quantiques,* J. Algebra **170** (1994), 229-265.

[2] N. Andruskiewitsch and F. Dumas, *On the automorphisms of $U_q^+(\mathfrak{g})$,* Preprint, 2003: http://arxiv.org/abs/math/0301066.

[3] K. A. Brown and K. R. Goodearl, *Lectures on algebraic quantum groups,* Advanced Courses in Mathematics, CRM Barcelona, Birkhäuser Verlag, Basel, 2002.

[4] G. Cauchon, *Effacement des dérivations et spectres premiers des algèbres quantiques,* J. Algebra **260** (2003), 476-518.

[5] G. Cauchon, *Spectre premier de $O_q(M_n(k)$: image canonique et séparation normale.* J. Algebra **260** (2003), 419-569.

[6] K. R. Goodearl and E. S. Letzter, *Prime factor algebras of the coordinate ring of quantum matrices,* Proc. Amer. Math. Soc. **121** (1994), 1017-1025.

[7] K. R. Goodearl and R. B. Warfield Jr., *An introduction to noncommutative nœtherian rings,* London Math. Soc. Student texts **16**, Cambridge University Press, Cambridge (1989).

[8] M. Gorelik, *The prime and primitive spectra of a quantum Bruhat cell translate,* J. Algebra **227** (2000), 211-253.

[9] J. E. Humphreys, *Introduction to Lie Algebras and Representations Theory,* Gratuate Texts in Mathematics **9**, Springer Verlag, New York (1980).

[10] T. J. Hodges and T. Levasseur and M. Toro, *Algebraic structure of multiparameter quantum groups,* Advances in Maths. **126** (1997), 52-92.

[11] J. C. Jantzen, *Lectures on quantum groups,* Graduate studies in mathematics **6,** Amer. Math. Soc., Providence (1996).

[12] A. Joseph, *Quantum Groups and Their Primitive Ideals,* Ergebnisse der Mathematik **(3) 29,** Springer-Verlag, Berlin (1995).

[13] S. Launois, *Combinatorics of $H-$ primes in quantum matrices,* J. Algebra **309** (2007), 139-167.

[14] S. Z. Levendorskii and Ya. S. Soibelman, *Algebras of functions on a compact quantum group, Schubert cells and quantum tori,* Comm. Math. Phys. **139** (1991), 141-170.

[15] T. Lam and L. Williams, *Total positivity for cominuscule grassmannians,* Preprint, 2007: http://arxiv.org/abs/math/0710.2932v1.

[16] R. Marsh and K. Rietsch, *Parametrisations of flag varieties,* Representation Theory, **8** (2004), 212-242.

[17] A. Mériaux, *Les diagrammes de Cauchon pour $U_q^+(\mathfrak{g})$,* Preprint, 2008: http://arxiv.org/abs/0807.1012

[18] A. Postnikov, *Total positivity, Grassmannians and Networks,* Preprint, 2006: http://arxiv.org/abs/math/0609764.

[19] H. Yamane, *A P.B.W. theorem for quantized universal envelopping algebra of type A_n*, Publ. Res. Inst. Math. Sci. **25** (1989), 503-520.